

Pub. Feb 96

11

PCT

WORLD INTELL



INTERNATIONAL APPLICATION PUBLISHED

NO 9604296A1

(51) International Patent Classification ⁶ :

C07J 7/00, 9/00, 13/00, 53/00, 71/00,
A61K 31/56, 31/57, 31/575, 31/58

(43) International Publication Date: 15 February 1996 (15.02.96)

(21) International Application Number: PCT/US95/09634

(22) International Filing Date: 4 August 1995 (04.08.95)

(30) Priority Data:

08/286,073 4 August 1994 (04.08.94) US
Not furnished 3 August 1995 (03.08.95) US

(71) Applicant: PHERIN CORPORATION [US/US]; Suite 240,
535 Middlefield Road, Menlo Park, CA 94025 (US).

(72) Inventors: JENNINGS-WHITE, Clive, L.; 3096 South 2300
East, Salt Lake City, UT 84109 (US). BERLINER, David,
L.; 380 Selby Lane, Atherton, CA 94027 (US). ADAMS,
Nathan, William; 4858 South Rockhill Lane, Salt Lake City,
UT 84123 (US).

(74) Agent: SUYAT, Reginald, J.; Fish & Richardson P.C., Suite
100, 2200 Sand Hill Road, Menlo Park, CA 94025 (US).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH,
CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG,
KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW,
MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT,
UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES,
FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD,
TG), ARIPO patent (KE, MW, SD, SZ, UG).

Published

With international search report.

(54) Title: PREGNANES AND CHOLANES AS NEUROCHEMICAL INITIATORS OF CHANGE IN HYPOTHALAMIC FUNCTION.

(57) Abstract

The invention relates to a method of altering hypothalamic function in an individual. The method comprises nasally administering a human vomeropherin, e.g. a pregnane or cholane steroid, or a pharmaceutical composition containing a vomeropherin, such that the vomeropherin binds to a specific neuroepithelial receptor. The steroid or steroids is/are preferably administered in the form of a pharmaceutical composition containing one or more pharmaceutically acceptable carriers. Other embodiments of the invention include pharmaceutical compositions containing the steroids.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

PREGNANES AND CHOLANES AS NEUROCHEMICAL INITIATORS OF CHANGE IN HYPOTHALMIC FUNCTION

5

Technical Field

This invention relates generally to pharmaceutical compositions and methods for effectuating change in human hypothalamic function, thereby altering certain behavior and physiology mediated by the hypothalamus of individuals. More particularly, the invention relates to the use of certain pregnane and cholane steroids as neurochemical effectuators of physiology and behavior.

Description of the Related Art

15 The present invention relates to certain compounds, namely pregnane and cholane steroids, and methods of using these compounds as human vomeropherins in order to alter hypothalamic function, thereby affecting certain consequent behavior and physiology, e.g., the reduction of anxiety. Pregnane steroids are characterized by a four ring steroidal structure, a methylation at the 13-position and at the 10-position, and ethylation at the 17-position. Pregnenes are a subset of pregnanes and have at least one double bond.

25 Ohloff, G. et al. (Helv. Chim. Acta (1983) 66:192-217), which is incorporated herein by reference, have shown that several steroids (androstenes) have an odor which varies with different isomeric, diastereomeric, and enantiomeric forms. Some members of this group have been reported to act as a pheromone in some mammalian species - for instance, 5 α -androst-16-en-3-one and 5 α -androst-16-en-3 α -ol in pigs (Melrose, D.R., et al., Br. vet. J. (1971) 127:497-502). These 16-androstenes produced by

30

- 2 -

the boar induce mating behavior in estrus sows (Claus, et al., Experimentia (1979) 35:1674-1675).

Cholane steroids are characterized by a 5-carbon side chain, the 2-pentyl group, at C-17 of the steroid nucleus. Some studies have noted that, in some species, various characteristics of certain 16-androstenes (including 5 α -Androst-16-en-3 α -ol and 5 α -Androst-16-en-3-one), such as concentration, metabolism, and localization, are sexually dimorphic (Brooksbank et al., J. Endocr. (1972) 52: 239-251; Claus, et al., J. Endocr. (1976) 68:483-484; Kwan, et al., Med. Sci. Res. (1987) 15:1443-1444). For instance, 5 α -Androst-16-en-3 α -ol and 5 α -Androst-16-en-3-one, as well as Androsta-4,16-dien-3-one, have been found at different concentrations in the peripheral blood, saliva and axillary secretions of men and of women (Kwan, T.K., et al., Med. Sci. Res. (1987) 15:1443-1444), and their function as a human pheromone, to the extent of affecting choice and judgment, has been suggested (Id.; see also Gower, et al., "The Significance of Odorous Steroids in Axillary Odour", In, Perfumery, pp. 68-72, Van Toller and Dodds, Eds., Chapman and Hall, 1988); Kirk-Smith, D.A., et al., Res. Comm. Psychol. Psychiat. Behav. (1978) 3:379). Androstenol (5 α -androst-16-en-3 α -ol) has been claimed to exhibit a pheromone-like activity in a commercial men's cologne and women's perfume (Andron™ for men and Andron™ for women by Jövan). Japanese Kokai No. 2295916, refers to perfume compositions containing androstenol and/or its analogues. Androstadien-3 β -ol (and perhaps the 3 α -ol) has also been identified in human axillary secretion (Gower, et al., supra, at 57-60). On the other hand, there is little agreement in the literature as to whether or not any putative pheromone actually plays any role in the sexual or reproductive behavior of mammals, particularly of humans. See:

- 3 -

Beauchamp, G.K., et al., "The Pheromone Concept in Mammalian Chemical Communication: A Critique", In: Mammalian Olfaction, Reproductive Processes and Behavior, Doty, R.L., Ed., Academic Press, 1976). See also: Gower, et al., supra at 68-73.

The pheromone properties of some estrene steroids for some mammalian species has been described. Michael, R.P. et al., Nature (1968) 218:746 refers to Estrogens (particularly Estradiol) as a pheromonal attractant of male rhesus monkeys. Parrot, R.F., Hormones and Behavior (1976) 7:207-215, reports Estradiol benzoate injection induces mating behavior in ovariectomized rats; and the role of the blood level of Estradiol in male sexual response (Phoenix, C.H., Physiol. and Behavior (1976) 16305-310) and female sexual response (Phoenix, C.H., Hormones and Behavior (1977) 8:356-362) in Rhesus monkeys has been described. On the other hand, there is little agreement in the literature as to whether or not pheromones as such play any role in the reproductive behavior and interpersonal communication of mammals (Beauchamp, G.K., et al., The Pheromone Concept in Mammalian Chemical Communication: A Critique', In: Mammalian Olfaction, Reproductive Processes, and Behavior, Doty R.L., Ed., Academic Press, 1976).

An embodiment of the subject invention concerns the non-systemic, nasal administration of certain pregnane and cholane steroids to affect a specific behavioral or physiological response in human subjects, e.g., a reduction of negative affect, mood, and character traits. In particular, nasal administration provides for contacting neurochemical receptors of a heretofore poorly understood neuroendocrine structure, commonly known as the vomeronasal organ ("VNO"; also known as "Jacobson's organ"), with one or more steroid(s) or with compositions containing the steroid(s). This organ is accessed

- 4 -

through the nostrils of most higher animals - from snakes to humans, and has been associated, inter alia, with pheromone reception in certain species (see generally Muller-Schwarze & Silverstein, Chemical Signals, Plenum Press, New York (1980)). The axons of the neuroepithelia of the vomeronasal organ, located supra palatal, form the vomeronasal nerve and have direct synaptic connection to the accessory olfactory bulb and indirect input from there to the cortico-medial amygdaloid basal forebrain and hypothalamic nuclei of the brain. The distal axons of terminalis nerve neurons may also serve as neurochemical receptors in the VNO. Stensaas, L.J., et al., J. Steroid Biochem. and Molec. Biol. (1991) 39:553. This nerve has direct synaptic connection with the hypothalamus.

Johnson, A. et al. (J. Otolaryngology (1985) 14:71-79) report evidence for the presence of the vomeronasal organ in most adult humans, but conclude that the organ is probably non-functional. Contravening results which suggest that the VNO is a functional chemosensory receptor are reported by Stensaas, L., et al., supra; and by Moran, D.T., et al., Garcia-Velasco, J. and M. Mondragon; Monti-Bloch, L. and B. Grosser all in J. Steroid Biochem. and Molec. Biol. (1991) 39.

It is apparent that it would be desirable to identify and synthesize human vomeropherins and pheromones and to develop pharmaceutical compositions and methods of use to influence hypothalamic function. This invention relates to the unexpected discovery that, when nasally administered to human subjects, certain neurochemical ligands, particularly pregnane and cholane steroids, and related compounds, or pharmaceutical compositions containing pregnanes, cholanes or related compounds, specifically bind to chemoreceptors of certain nasal neuroepithelial cells and this binding generates a

- 5 -

series of neurophysiological responses resulting in an alteration of hypothalamic function of an individual. When properly administered, the effect of certain of these compounds on the hypothalamus affects the function of the autonomic nervous system and a variety of behavioral-or physiological phenomena which include, but are not limited to the following: anxiety, premenstrual stress, fear, aggression, hunger, blood pressure, and other behavioral and physiological functions normally regulated by the hypothalamus. See Otto Appenzeller, The Autonomic Nervous System. An Introduction of basic and clinical concepts (1990); Korner, P.I. Central nervous control of autonomic cardiovascular function, and Levy, N.M. and Martin, P.J. Neural control of the heart, both in Handbook of Physiology; Section 2; Cardiovascular System - the heart, Vol. I, Washington DC, 1979, American Physiological Society; Fishman, A.P., et al. editors, Handbook of Physiology. Section 3: Respiratory System. Vol. II. Control of breathing, Bethesda MD, 1986. American Physiological Society.

In some instances a single pregnane or cholane steroid, or related compound, is administered, in some instances combinations of pregnane and/or cholane steroids and/or related compounds are administered and in some instances one or more pregnane or cholane steroids are co-administered along with one or more estrane or estrene steroids, androstane or androstene steroids or a related compound.

Summary of the Invention

Accordingly, it is an object of this invention to provide pharmaceutical compositions which contain human vomeropherins or pheromones and are suitable for nasal administration in an individual.

- 6 -

It is also an object of this invention to provide methods of using these compositions to alter hypothalamic function of an individual.


It is a further object of this invention to
5 provide methods of using these compositions to affect physiological and behavioral functions of individuals which are normally regulated by the hypothalamus.

Finally, it is an object of this invention to provide methods of altering hypothalamic function which
10 have the following advantages: 1) administration directly to the chemoreceptors in the nasal passage and the vomeronasal organ, without pills or needles - i.e., non-invasively; 2) a mode of drug action through the invention system and not through the circulatory system -
15 thus brain function can be affected without consideration of the bloodbrain barrier; 3) a direct means of affecting the hypothalamus - there is only one synaptic junction between pheromone receptors and the hypothalamus; and,
4) providing a highly specific drug effect, thereby
20 greatly reducing the potential for undesirable side-effects - this because sensory nerves are addressed to a specific location in the brain.

Additional objects, advantages and novel features of the invention will be set forth in part in the
25 description which follows, and in part will become apparent to those skilled in the art upon examination of the following, or may be learned by practice of the invention.

Objects of this invention are achieved by
30 providing a pharmaceutical composition suitable for nasal administration in an individual. The composition contains a pharmaceutically acceptable carrier and a pregnane steroid with the formula:

- 7 -



wherein P_1 is selected from the group consisting of oxo, α -(β -) hydroxy, α -(β -) acetoxy, α -(β -) propionyloxy, α -(β -) methoxy, α -(β -) lower acyloxy, α -(β -) lower alkyloxy, and α -(β -) benzoyloxy; P_2 is selected from the group
 5 consisting of methyl, hydroxymethyl, acyloxymethyl, alkoxyethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxyalkyl; P_3 is selected from the group consisting of hydrogen, oxo, halo, hydroxy, alkoxy, and acyloxy; P_4 through P_{12} may each be, independently, hydrogen, halo,
 10 methyl, or halo-, dihalo-, or perhalomethyl and when P_2 is methyl and P_3 is β -hydroxy, P_2 and P_3 may be jointed to form a cyclic ether; P_{13} is hydrogen, methyl, methylene, halo-substituted methyl, halo-substituted methylene, ethyl, ethylenyl, acetylenyl, methyl-methylenyl, methyl-
 15 methinyl; and "a", "b", "c", "d", "e", "h", "i", and "j" are alternative sites for optional double bonds, and "j" or "k" may also be triple bonds. Halo substituents include fluoro, bromo, chloro and iodo atoms.


One class of preferred steroids has "b" as a
 20 double bond, particularly wherein "d" or "e" is also a double bond. Another preferred class has "a" and "c" as double bonds, or only "c" as a double bond. Yet another preferred class contains "h" as a double bond, with i and j being absent (i.e., single bonds), j being a double

- 8 -

bond, or j being a triple bond. In another class, "h" is absent, and j or i is a double bond, or i and j are absent, or j and i are double bonds, or j is a triple bond.

- 5 The term lower alkyl, lower alkoxy, etc., is meant to encompass carbon chains of 1 to 6 carbon atoms, preferably 1 to 4 carbon atoms.

The invention also is directed to novel pregnane and cholane derivations of the formula



- 10 wherein P₁ is selected from the group consisting of oxo, α-(β-) hydroxy, α-(β-) acetoxy, α-(β-) propionyloxy, α-(β-) methoxy, α-(β-) lower acyloxy, α-(β-) lower alkyloxy, and α-(β-) benzyloxy; P₂ is selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl, 15 alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxylalkyl; P₃ is selected from the group consisting of hydrogen, oxo, halo, hydroxy, alkoxy, and acyloxy; P₄ through P₁₂ may each be, independently, hydrogen, halo, methyl, or halo-, dihalo- or perhalomethyl; P₁₃ is 20 hydrogen, methyl, methylene, halo-substituted methyl or halo-substituted methylene, ethyl, ethylenyl, acetylenyl, methyl-methylenyl or methyl-methinyl; and "a", "b", "c", "d", "e", "h", "i", "j" and "k" are alternative sites for optional double bonds and "j" or "k" may also be triple 25 bonds; and when P₂ is methyl and P₃ is β-hydroxy, P₂ and P₃

may be joined to form a cyclic ether; with the provisos that

- i) when $P_1 = \text{oxo}$; b is present; e, a, c, d are absent; $P_2 = \text{methyl}$; and $P_3, P_4, P_7, P_9, P_{10}, P_{11}$ and P_{13} are hydrogen; then if h is present or absent, P_6 must be present and cannot be hydrogen;

- ii) when $P_1 = \text{OH}$ and c is present; e, a, b and d are absent; $P_2 = \text{methyl}$; and $P_3, P_4, P_7, P_9, P_{10}, P_{11}, P_{12}$ and P_{13} are hydrogen; then
- (a) if h is present, P_6 must be present and is not hydrogen;
 - (b) if h is absent and P_6 is present, P_6 cannot be hydrogen;
 - (c) if h is absent and P_6 is present, then i and j are both present;

- iii) when $P_1 = \beta\text{-OH}$ and b is present; e, a, c, d are absent; $P_2 = \text{methyl}$; and $P_3, P_4, P_7, P_9, P_{10}, P_{11}$ and P_{13} are hydrogen;
- (a) if h is present, i and/or j must be present; or P_6 is not hydrogen;
- (b) if h is present, j cannot be a triple bond;

- iv) when $P_1 = \alpha\text{-OH}$; b is present; e, a, c, d are absent; $P_2 = \text{methyl}$;
and $P_3, P_4, P_7, P_9, P_{10}, P_{11}$ and P_{13} are hydrogen; then
(a) if h is present, i must be absent
and P_6 cannot be hydrogen;

- 10 -

- (b) if i and j are both absent, P₆ cannot be hydrogen;
- v) when P₁ = oxo; and b and d are present; e, a, and c are absent; P₂ = methyl; and P₃, P₄, P₇, P₉, P₁₁ and P₁₃ are all hydrogen; then if h is absent and i is present, P₆ must be present and cannot be hydrogen;
- vi) when P₁ and P₃ are oxo and b is present; h, e, a, c and d are absent; P₂ = methyl; P₄, P₅, P₇, P₈, P₉, P₁₀, P₁₁ and P₁₃ are hydrogen; then
- (a) if P₆ is absent, j cannot be a double bond;
- (b) if P₆ is hydrogen, i or j must be present;
- vii) when P₁ = OMethyl, a and c are present; and P₂ = methyl; e, b, d and h are absent and
- P₃, P₄, P₇, P₈, P₉, P₁₀, P₁₁ and P₁₃ are hydrogen; then
- (a) if P₆ is hydrogen, i or j must be present;
- (b) if P₆ is absent, i and j cannot both be double bonds;
- viii) when P₁ = oxo and a, b, c, d, e are absent;
- P₂ = methyl; and
- P₃, P₄, P₇, P₉, P₁₀, P₁₁, P₁₂ and P₁₃ are hydrogen; then

- 11 -

- (a) if h is absent and j is present as a double bond; P_6 cannot be hydrogen;
- (b) if h, i and j are absent, P_6 cannot be hydrogen; and
- (c) if h is present, j cannot be a triple bond;

ix) when $P_1 = \text{OH}$, a, b, c, d, e and h are absent;

$P_2 = \text{methyl}$;

$P_3, P_4, P_7, P_8, P_9, P_{10}, P_{11}, P_{12}$ and P_{13} are hydrogen; then

- (a) if j is a double bond and i is absent; then P_6 is not hydrogen;
- (b) if i is a double bond and j is absent, then P_6 is not hydrogen;
- (c) i and j cannot both be double bonds;
- (d) if i and j are absent, P_6 cannot be hydrogen;

x) when $P_1 = \text{oxo}$, e and b are present; and a, c, d are absent;

$P_2 = \text{methyl}$; and

$P_3, P_4, P_7, P_9, P_{10}, P_{11}$ and P_{13} are hydrogen; then

- (a) if h is present, j cannot be a triple bond;
- (b) if h is absent and i is a double bond, then P_6 is not hydrogen;
- (c) if h is absent and i is a double bond, then P_6 is not hydrogen;

- 12 -

(d) if h is absent and neither i nor j is a double bond, then P₆ is not hydrogen;

5 xi) when P₁ = oxo, P₃ = OH, b is present, a, e, c, d, h, i, j are absent, P₂ = methyl and P₄, P₇, P₉, P₁₀, P₁₁ and P₁₃ are hydrogen; then P₆ cannot be hydrogen;

xii) when P₁ = oxo, b is present, a, e, c, d, h, i are absent, P₆ and P₂ are methyl and

10 P₃, P₄, P₅, P₇, P₈, P₉, P₁₀ and P₁₁ are hydrogen; then

(a) if j is a double bond, P₁₃ cannot be ethylenyl;

15 (b) if neither j nor k is a double or triple bond, then P₁₃ cannot be acetylenyl;

xiii) when P₁ = OH, c is present and a, b, e, d, h and i are absent;

P₂ and P₆ are methyl; and

20 P₃, P₄, P₅, P₇, P₈, P₉, P₁₀, P₁₁ and P₁₂ are hydrogen; then

(a) if j is a double bond, P₁₃ is not ethyl;

25 (b) if j is absent and k is a double bond, P₁₃ is not methyl-methylenyl;

- 13 -

- (c) if j and k are absent; P_{13} is not ethyl; and
- (d) if k is a triple bond, P_{13} cannot be methyl-methinyl.

5 Other objects of this invention are achieved by providing a method of altering hypothalamic function and/or autonomic function in an individual. A ligand for a chemoreceptor displayed on the surface of a nasal neuroepithelial cell is provided wherein the cell is a
10 part of tissue other than olfactory epithelia; and, the ligand is administered within a nasal passage of the individual such that the ligand binds specifically to the chemoreceptor, resulting in an alteration of hypothalamic function of the individual.

15 All embodiments of this application relate to and include the functional equivalents of the steroid structures disclosed in these embodiments and to those modified steroids which demonstrate said functional equivalence, whether or not the modified steroids are
20 explicitly disclosed.

Brief Description of the Drawings

FIG. 1 is the data on the integrated EVG, GSR and ST for compound A1-P1 in males as tested according to Examples 16 and 17.

25 FIG. 2 is the data for the integrated EVG for compounds A1-P1, A2-P1, A4-P1, A3-P1, A1-P4, A2-P4 in females.

FIG. 3 is the data for the ST measurements of compounds A1-P1, A2-P1, A4-P1, A3-P1, A1-P4, A2-P4 in
30 females.

- 14 -

FIG. 4 is the data for the GSR measurements in females of compounds A1-P1, A2-P1, A4-P1, A3-P1, A1-P4, A2-P4.

FIG. 5 is the data for ST, GSR and EVG
5 measurements in females of compound A1-P3.

FIG. 6 is the data for RF and EKG measurements in females of compound A1-P3.

FIG. 7 is the data for EEG measurements in females of the compound A1-P3.

10 FIG. 8 is the data for the ST, GSR and EVG measurements in males of compound A1-P3.

FIG. 9 is the data for the RF and EKG measurements in males of compound A1-P3.

FIG. 10 is the data for the EEG measurements in
15 males of compound A1-P3.

FIGS. 11 and 12 show the data of the ST, GSR and EVG measurements in males and females, respectively, for compound A2-P3.

FIGS. 13 and 14 show the data of the EEG
20 measurements in males and females, respectively, for compound A2-P3.

FIGS. 15 and 16 show the data for the RF and EKG measurements in males and females, respectively, for compound A2-P3.

25 FIGS. 17 and 18 show the data of the ST, GSR and EVG measurements in males and females, respectively, for compound A8-P1.

FIGS. 19 and 20 show the data of the RF and EKG measurements in males and females, respectively, for
30 compound A8-P1.

FIGS. 21 and 22 show the data for EEG measurements in males and females, respectively, for compound A8-P1.

FIGS. 23 and 24 show the data for ST, GSR and EVG measurements in males and females, respectively, for
35 compounds A6-P1.

- 15 -

FIGS. 25 and 26 show the data for RF and EKG measurements in males and females, respectively, for compound A6-P1.

FIGS. 27 and 28 show the data for EEG measurements
5 in males and females, respectively, for compound A6-P1.

FIGS. 29, 30 and 31 show the data for ST, GSR, EVG, RF EKG and EEG measurements in males of 20,21-dimethylpregna-5,20-dien-3 β -ol.

FIGS. 32, 33 and 34 show the data for the ST, GSR,
10 EVG, RF KEG and EEG measurements in females of 20,21-dimethylpregna-5,20-dien-3 β -ol.

FIGS. 35, 36 and 37 show the data for the ST, GSR, EVG, RF KEG and EEG measurements in males of 20,21-dimethylpregna-5,20-dien-3-one.

FIGS. 38, 39 and 40 show the ST, GSR, EVG, RF EKG
15 and EEG measurements in females of 20,21-dimethylpregna-5,20-dien-3-one.

FIGS. 41, 42 and 43 show the ST, GSR, EVG, RF EKG and EEG measurements in males of compound A14-P2.

FIGS. 44, 45 and 46 show the ST, GSR, EVG, RF, EKG
20 and EEG measurements in females of compound A14-P2.

FIGS. 47, 48 and 49 show the ST, GSR, EVG, RF, EKG and EEG measurements in males of compound A7-P2.

FIGS. 50, 51 and 52 show the ST, GSR, EVG, RF, EKG
25 and EEG measurements in females of compound A7-P2.

FIGS. 53 and 54 show the ST, GSR, EVG, EEG measurements in males of compound A11-P1.

FIGS. 55 shows the data of EEG measurements in males of compound A13-P1.

FIGS. 56, 57 and 58 show the data of measurements
30 of ST, GSR, EVG, RF, EKG and EEG in females of compound A13-P1.

FIGS. 59 shows EVG, EDA and BT data of measurements in women of compound A3/P1.

- 16 -

FIG. 60 shows EVG, EDA and BT data of measurements in women of compound A4/P1.

FIGS. 61 and 62 show data of measurements in men and women, respectively, of compound A8/P1.

5 FIGS. 63 and 64 show data of measurements in men and women, respectively, of compound A13/P8.

FIGS. 65 and 66 show data of measurements in men and women, respectively, of compound A6/P1.

10 FIGS. 67 and 68 show data of measurements in men and women, respectively, of the 20-methyl derivative of compound A6/P1.

FIGS. 69 and 70 show data of measurements in men and women, respectively, of the 20,21-dimethyl derivative of compound A1/P1.

15 FIGS. 71 and 72 show data of measurements in men and women, respectively, of the 20,21-dimethyl derivative of compound A6/P1.

FIGS. 73 and 74 show the data of measurements in men and women, respectively, of compound A14/P2.

20 FIGS. 75 and 76 show the data of measurements in men and women, respectively, of compound A12/P1.

FIGS. 77 and 78 show the data of measurements in men and women, respectively, of compound A7/P2.

25 FIGS. 79 and 80 show the data of measurements in men and women, respectively, of compound A13/P1.

FIGS. 81 and 82 show the data of measurements in men and women, respectively, of compound A2/P7.

FIGS. 83 and 84 show the data of measurements in men and women, respectively, of compound A3/P5.

30 FIGS. 85-96 refer to the cholanes on Chart II.

FIGS. 85 and 86 show the data of measurements in men and women, respectively, of compound A8/C1.

FIGS. 87 and 88 show the data of measurements in men and women, respectively, of compound A2/C1.

- 17 -

FIGS. 89 and 90 show the data of measurements in men and women, respectively, of the acetate of compound A2/C1.

FIGS. 91 and 92 show the data of measurements in
5 men and women, respectively, of compound A1/C1.

FIGS. 93 and 94 show the data of measurements in men and women, respectively, of compound A3/C1.

FIGS. 95 and 96 show the data of measurements in men and women, respectively, of compound A13/C1.

10 Detailed Description of the Invention

I. Definitions

An "affect" is a transient feeling state. Typical negative affects are feelings of nervousness, tenseness, shame, anxiousness, irritability, anger, rage and the
15 like. "Moods" are longer lasting feeling states such as guilt, sadness, hopelessness, worthlessness, remorsefulness, misery, unhappiness and the like. "Character traits" are more permanent aspects of an individual's personality. Typical negative character
20 traits are sensitivity, regretfulness, blameworthiness, stubbornness, resentment, bitterness, timidity, laziness and the like.

The vomeropherins according to the present invention may have use for stimulating, through contact
25 with the VNO, of one or more of the hormonal, behavioral and autonomic functions of the hypothalamus. Due to the predominant role played by the hypothalamus in a wide variety of internal body functions and the neural connection between the VNO and the hypothalamus, the
30 vomeropherins according to the present invention are in a position to stimulate such functions as endocrine output control, for example, the control of the pituitary output of vasopressin and oxytocin as well as a number of other peptides. Vasopressin is an anti-diuretic hormone

- 18 -

because of its action within the kidney to enhance water uptake and to concentrate the urine. In addition, it has an action within the body to regulate blood pressure through its action on arterial smooth muscle and an
5 action on metabolism through its enhancement of glycogen conversion to glucose in the liver. Oxytocin, receptors of which are found on uterine smooth muscle and on mammary smooth muscle, can cause milk letdown via contraction of the mammary smooth muscle and cause
10 uterine contractions during birth. The hypothalamus also controls release of hormones from the anterior pituitary gland such as ACTH, prolactin, LH (Luteinizing Hormone), GH (Growth Hormone), TSH (Thyroid Stimulating Hormone), FSH (Follicle Stimulating Hormone) and beta-endorphin.
15 Thus, for example, the ability to control LH secretion may lead to control of fertility, in females, or testosterone production in males. Testosterone production may be utilized for treatment of conditions such as low libido in males or for treatment of muscle
20 wasting diseases or conditions, such as aging. Testosterone reduction may be utilized for treatment of conditions such as prostate cancer.

Control of behavior hypothalamic functions is also feasible by use of the vomeropherins according to the
25 present invention. It is known that the hypothalamus controls such behavioral outputs as fear, rage, pleasure and circadian rhythms which regulate sleep and wakefulness. Other functions controlled by the hypothalamus include appetite, thirst, sympathetic
30 functions such as, flight and fight, and functions such as cardiovascular control, thermoregulation and visceral functions such as control of the gut muscle and acid secretion for digestion. Thus, while there is a multitude of sensory inputs into the hypothalamus from
35 various parts of the anatomy it is believed that the

- 19 -

vomeropherins of the present invention provide, for the first time, a way of stimulating through the nasal cavity by inhalation to contact the epithelial cells in the VNO, a method of stimulating functions of the hypothalamus
5 discussed above.

"Pregnane steroids" are aliphatic polycyclic hydrocarbons characterized by a four-ring steroidal structure with a methylation at the 10- and 13-positions and ethylation (including unsaturated groups) at the 17-
10 position. A pregnane is a subset of pregnanes commonly understood to mean that the compound has at least one double bond. Furthermore, all derivatives which have the structural characteristics described above are also referred to generically as pregnane steroids.

15 A "cholane steroid" is an aliphatic polycyclic hydrocarbon characterized by a four-ring steroid structure, with methylation at the 10- and 13-positions and a 2-pentyl group (including unsaturated groups) at the 17-position.


20 A "chemoreceptor" is a receptor molecule displayed on the surface of a "chemosensory" neuroepithelial cell which binds in a stereospecific fashion to a particular ligand or ligands. This specific binding initiates a signal transduction which initiates an afferent nerve
25 impulse. Chemoreceptors are found, inter alia, in taste buds, olfactory epithelium and vomeronasal tissue.

"Pregnene steroids", as the term is used herein, are aliphatic polycyclic hydrocarbons with a four-ring steroidal structure, at least one double bond in the A-
30 ring, methylation at the 10-position and 13-position, ethylation (including unsaturated groups) at the 17-position and an oxo, hydroxyl or hydroxyl derivative such as alkoxy, ester, benzoate, cypionate, sulfate or glucuronide, at the 3-position. Derivatives which

- 20 -

contain these structural characteristics are also referred to generically as pregnene steroids.

The following structure shows the four-ring steroidal structure common to pregnane and pregnene steroids. In describing the location of groups and substituents, the following numbering system will be employed:



"Sexually dimorphic" refers to a difference in the effect of, or response to, a pharmaceutical agent between males and females of the same species.

An "effective amount" of a drug is a range of quantity and/or concentration which brings about a desired physiological and/or psychological effect when administered to an individual in need of the drug. In the present case, a needy individual is one with a physiological or behavioral trait which is normally regulated by the hypothalamus and wherein it is desirable to affect the function of the hypothalamus or the trait. The effective amount of a given drug may vary depending upon the function to be affected, the desired effect, route of administration, and the like. For example, when the steroid is administered as a solution applied to the facial skin of a subject an effective concentration is from 1 microgram/ml to 100 μ g/ml, preferably 10 to 50 μ g/ml and most preferably 20 to 30 μ g/ml. When the steroid is introduced directly into the VNO an effective amount is about 1 picogram to about 1 nanogram, more preferably about 10 picograms to about 50 picograms.

- 21 -

When the steroid is administered to the nasal passage, by ointment, cream or aerosol, or the like, an effective amount is about 100 pg to about 100 micrograms, preferably about 1 ng to about 10 micrograms. It follows that some drugs may be effective when administered by some routes, but not effective when administered by other routes.

The "hypothalamus" is the portion of the diencephalon comprising the ventral wall of the third ventricle below the hypothalamic sulcus and including structures forming the ventricle floor, including the optic chiasma, tuber cinereum, infundibulum, and mammillary bodies. The hypothalamus regulates the autonomic nervous system and controls several physiological and behavioral functions such as the so-called fight and flight responses, sexual motivation, water balance, sugar and fat metabolism, hunger, regulation of body temperature, endocrine secretions, and others. The hypothalamus is also the source of vasopressin which regulates blood pressure, and oxytocin which induces parturition and milk release. All hypothalamic functions are potentially modulatable by the vomeropherin therapy described herein.

A "ligand", as used herein, is a molecule which acts as a chemical signal by specifically binding to a receptor molecule displayed on the surface of a receptor cell, thereby initiating a signal transduction across the cell surface. Binding of ligands to chemosensory receptors can be measured. Chemosensory tissue, such as vomeronasal neuroepithelium or olfactory neuroepithelium, contains a multiplicity of neuroreceptors cells, each displaying at least one cell surface receptor. Many of the receptor molecules have identical ligand specificity. Therefore, when the tissue is exposed to a ligand for which it has specificity (for example a exposure of the

- 22 -

VNO to a vomeropherin) a summated change in cell surface receptor potential can be measured.

As used herein, "lower alkyl" means a branched or unbranched saturated hydrocarbon chain of 1 to 4 carbons, such as, for example, methyl, ethyl, n-propyl, i-butyl and the like. "Alkoxy" as used herein is used in its conventional sense to mean the group -OR wherein R is alkyl as herein defined.

A "pheromone" is a substance that provides chemical means of communication between members of the same species through secretion and peripheral chemoreception. In mammals pheromones are usually detected by receptors in the vomeronasal organ of the nose. Commonly, pheromones effect development, reproduction and related behaviors. A "vomeropherin" is a more general term which includes pheromones and describes a substance from any source which functions as a chemosensory messenger, binds to a specific vomeronasal neuroepithelial receptor, and induces a physiological or behavioral effect. The physiologic effect of a "vomeropherin" is mediated through the vomeronasal organ.

A picogram (pg) is equal to .001 nanograms (ng). A ng is equal to .001 micrograms (μ g). A μ g is equal to .001 mg.

- 23 -


















































































































II. Modes for Carrying Out the InventionA. Pregnanes useful in the Invention

The invention is directed to a group of certain pregnane steroids.

5 A subset of pregnanes within the group are believed to be novel. Syntheses are described herein for the following compounds as designated on the chart:

Chart 1 includes pregnanes to which the invention is directed, but do not limit its scope. The synthesis
10 diagrams that follow depict intermediate and substructure syntheses for the preparation of these pregnanes:

PREGNANCE

	1	2	3	4	5	6	7	8
1								
2								
3								
4								
5								
6								
7								
8								
9								
10								
11								
12								
13								
14								


SUBSTRUCTURE SYNTHESSES

Referring to the preceding table, the following are exemplary syntheses for intermediates in a given row (A1 through A13) or column (P1 through P8).

5


SUBSTRUCTURE SYNTHESSES: TYPE A

A1:






Percy L. Julian, Edwin W. Meyer and Helen C. Printy,
J. Amer. Chem. Soc., 1948, 70, 3, 887.

Also a commercially available substructure, for
10 example, 17 α -ethynyltestosterone.









- 26 -

A2:




This is a commercially available substructure, for example, dehydroepiandrosterone, pregnenolone.

A3:




- 5 David G. Loughhead, *J. Org. Chem.*, 1985, Vol. 50, No. 20, p. 3931.

A4:



I. Z. Kabore, Q. Khuong-Huu, and A. Pancrazi,
Tetrahedron, 1978, Vol. 34, p. 2807.


A5:




I. Dory, G. Szabo and P. Opoczky, *Acta Chim, Hung.*,
Vol. 20, p. 67 (1959).

Bernhard Krieger, Egbert Blanke, and Emanuel Kaspar,
5 German Patent 1,297,603 (1969).

A6:




Alan M. Krubiner, Norman Gottfried, and Eugene P. Oliveto, *J. Org. Chem.*, 1969, **34**, 11, 3502.




Roberto Sciaky and Alberto Consonni, *Gazz. Chim. Ital.*, 1962, **92**, 730.

- 30 -

A7:



A8:




See example.




Vladimir Petrow, Yueh-sha Wang, Leon Lack, Avery
 5 Sandberg, Nobuyuki Kadohama, and Keith Kendle, *J.*
Steroid Biochem., 1983, 19, 1491.

A9:




Steven R. Schow and Trevor C. McMorris, *Steroids*,
1977, Vol. 30, No. 3, p. 389.

Also a commercially available substructure, for
5 example, 17 α -ethynyldihydrotestosterone.




A10:




This is a commercially available substructure, for example, pregnanolone, androsterone.

Also:




- 5 J.M. Kohli, A Zaman and A.R. Kidwai, *Phytochemistry*, 1971, Vol. 10, p. 442.

A11:




See example.





Frederick Brown and Carl Djerassi, *J. Amer. Chem. Soc.*, 1980, 102, 2, 807.



S A12:



A13:




MCPBA


MCPBA
24%

SUBSTRUCTURE SYNTHESSES: TYPE P

P1:




Ajay K. Bose and N.G. Steinberg, *Synthesis*, 1970, p. 595.




5 Steven R. Schow and Trevor C. McMorris, *Steroids*, 1977, vol. 30, No. 3, p. 389.

P2:




Ronald Breslow and Louis M. Maresca, *Tetrahedron Letters*, 1977, No. 7, p. 623.




Braja G. Hazra, Vandana S. Pore, Padmakar L. Joshi,
5 *J. Chem. Soc., Perkin Trans I*, 1993, (15), 1819-22.

Also a commercially available substructure, for
example, 5 α -pregn-17(20)-en 3 β -ol (Steraloids):




P3:




This is a commercially available substructure,
for example, pregna-5, 16-dien-3 β -ol (Steraloids).

When commercially unavailable, synthesis
5 proceeds as below:




John P. Dusza and Werner Bergman, *J. Org. Chem.*,
1960, 25, 79.

P4:




C.W. Shoppee, Ruth E. Lack, and B.C. Newman,, J.
Chem. Soc., 1964, p. 3188.

Also a commercially available substructure, for
5 example, 5 α -pregnan-3 β -ol (Steraloids):




- 39 -

P5:




Alan M. Krubiner, Norman Gottfried, and Eugene P. Oliveto, *J. Org. Chem.*, 1969, Vol. 34, No. 11, p. 3502.

5 P6:




Eugene P. Oliveto, Corrine Gerold, and Lois Johnson, *J. Am. Chem. Soc.*, 1951, 73, 5073.

P7:




Pierre Crabble and Esperanza Velarde, U.S. Patent
3,681,410, 1972.

P8:




Maya Dvolaitzky Anne M. Giroud, and Jean Jacques,
Bull. Soc. Chim. France, 1963, 62.



French Patent 1,536,034, 1968.

17 α -PREGNANES


For substructures P1, P4, and P5, the normal configuration at the 17-position is β . However, the corresponding 17 α analog may also be prepared by using 17 α -pregnolone as the starting material. For example:




Alan M. Krubiner, Norman Gottfried, and Eugene P. Oliveto, *J. Org. Chem.*, 1969, Vol. 34, No. 11, p. 3502.

METHYLPREGNANES


The following methodology enables a methyl group to be placed at the 20-position whenever allowed by the structure, namely with P1, P2, P3, P4 and P6:



- 5 J. Bryan Jones and Keith D. Gordon, *Can. J. Chem.*, 1972, vol. 50, p. 2712.




John P. Dusza and Werner Bergmann, *J. Org. Chem.*,
1960, 25, 79.




David G. Loughhead, J. Org. Chem., 1985, 50, 3931.


U.S. Patent 3,681,410 teaches preparation of 6α-methyl analogs:




U.S. Patent 3,492,318 teaches preparation of 18- and 21-methyl analogs:



Certain methylated pregnenolone precursors are commercially available, viz 6, 16 α (β)-methyl:



In addition 17 α -methylpregnenolone is readily available: French patent 1,363,191:




Therefore, compounds synthesized from pregnenolone may also be prepared with methyl groups at the 6, 16, or 17 positions by using the appropriate methylpregnenolone precursor.

- 5 Dimethyl compounds, such as the described 18, 21-dimethylpregna-4,16-dien-20-yn-3-one, may be prepared by one of three general methods:

The first method combines a methylated precursor, such as those in the 6, 16, or 17-positions, with
10 methodology which introduces a methyl group, such as in the 20-position.

The second method uses a dimethylated precursor, such as the commercially available 6, 16 α - dimethylpregnenolone.

- 15 The syntheses of other dimethylated pregnenolone precursors have been described, as in the following examples:




Sylvestre Julia, Colette Neuville, and Pierre Simon,
Bull. Soc. Chim. France, 1962, 1495.


- 48 -



Elliot Shapiro, Theodore Legatt, Lois Weber, Merl Steinberg, A. Watnick, M. Eisler, Marilyn Gilmore Hennessey, C.T. Coniglio, W. Charney, and Eugene P. Oliveto, *J. Med. Pharm. Chem.* 1962, 5, 975.




5 James Cairns, Colin L. Hewett, Robert T. Logan, George McGarry, Donald F. M. Stevenson, and Gilbert F. Woods, *J.C.S. Perkin I.* 1976, 1558.




R. Deghenghi and R. Gaudry, *J. Amer. Chem. Soc.*, 1961, 4668.

British Patent 927,515:




Romano Deghenghi and Roger Gaudry, *Tetrahedron Letters*, 1962, No. 11, p. 489.




- 50 -

W. J. Adams, D.K. Patel, V. Petrov, I. A. Stuart-Webb, and B. Sturgeon. J. Chem. Soc., 1956, 4490.

The third method starts with an unmethylated precursor, such as pregnenolone, and utilizes
5 methodology which introduces two methyl groups, as in the following example:





The 20, 21-dimethyl pregnanes are also known as 24-norcholanes. 24-norcholanes may alternatively be prepared by degradation of a cholane precursor, as in
10 the following example:




Yutaka Hirano, Tadashi Eguchi, Masaji Ishiguro, and Nobuo Ikekawa. Chem. Pharm. Bull., 1981, 31(2), 394.

HALOPREGNANES

U.S. Patent 3,681,410 teaches the preparation of:




Derek H. R. Barton, George Bashlars and Jean-Louis
5 Fourrey, Tetrahedron Letters, 1983, vol. 24, 1605.




Biao Jiang and Yuanyao Xu, *Tetrahedron Letters*, 1992,
vol. 33, 511.

Certain methylated pregnenolone precursors are commercially available, viz 6, 16 α (β)-methyl:




















































In addition 17 α -methylpregnenolone is readily available: French patent 1,363,191:



Therefore, compounds synthesized from pregnenolone may also be prepared with methyl groups at the 6, 16, or 17 positions by using the appropriate methylpregnenolone precursor.

CHART II

CHOLANES


<u>A</u> 9	1	2	3	4	5	6	7
1	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL
2	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL
3	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL
6	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL
8	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL
11	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL
13	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL	 NOVEL

SUBSTRUCTURE SYNTHESSES

Referring to the preceding table, the following are exemplary syntheses for the substructures in a given row (A1 through A13) or column (C1 through C7).


SUBSTRUCTURE SYNTHESSES: TYPE A

A1:




See Example.


A2:




This is a commercially available substructure, for example PREGNENOLONE, STIGMASTEROL, CHOLENIC ACID.

A3:


See Example.

A6:


See Example.

A8:


See Example.

A11:


D. H. R. Barton, J. Boivin, D. Crich, and C. H. Hill, *J. Chem. Soc. Perkin Trans. I*, 1986, p. 1805.

A13:

See Example.


SUBSTRUCTURE SYNTHESSES: TYPE CC1:

J. P. Schmit, M. Piraux, and J. F. Pilette, *J. Org. Chem.*, 1975, Vol. 40, No. 11, p. 1586.


C2:

W. Bergmann and J. P. Dusza, *J. Org. Chem.*, 1958, Vol. 23, p. 1245.



- 60 -





J. E. van Lier and L. L. Smith, *J. Org. Chem.*, 1970, Vol. 36, No. 8, p. 2631.



A. Burger, F. Colobert, C. Hetru, and B. Luu, *Tetrahedron*, 1988, Vol. 44, No. 4, p. 1141.

C5:A. B. Turner, *Chemistry and Industry*, 1979, p. 385.**C6:**


G. Stoeck and H. Stein, German Patent No. 854,517, 1952.

C7:


A. Burger, J-P. Roussel, C. Hetru, J. A. Hoffmann, and B. Luu,
Tetrahedron, 1989, Vol. 45, No. 1, p. 155.

24-NORCHOLANES and 24-METHYLCHOLANES

Structures with the side chain shortened or lengthened by one carbon atom at the 24-position may be prepared using analogous methodology. 23-Methylcholanes are also available by similar means. Examples follow.




23-METHYL




24-METHYL

J. P. Schmit, M. Piraux, and J. F. Pilette, *J. Org. Chem.*, 1975, Vol. 40, No. 11, p. 1586.




24-METHYL


Y. M. Sheikh and C. Djerassi, *Steroids*, 1975, Vol. 26(1), p. 129.



24-NOR




23-METHYL





24-METHYL

M. Morisaki, M. Shibata, C. Duque, N. Imamura, and N. Ikekawa, *Chem. Pharm. Bull.*, 1980, Vol. 28(2), p. 606.




D. H. R. Barton, J. Boivin, D. Crich, and C. H. Hill, *J. Chem. Soc. Perkin Trans. I*, 1986, p. 1805.



A. Burger, F. Colobert, C. Hetru, and B. Luu, *Tetrahedron*, 1988, Vol. 44, No. 4, p. 1141.



B. M. Trost, R. J. Kulawiec, and A. Hammes, *Tetrahedron Letters*, Vol. 34, No. 4, p. 587.



A. Burger, J-P. Roussel, C. Hetru, J. A. Moffmann, and B. Luu, *Tetrahedron*, 1989, Vol. 45, No. 1, p. 155.

- 65 -

C. Synthetic Methods

1. Preparation of 3-, 6-, 19-, 20- and 21-position derivatives.

The compounds used in the methods of this invention are pregnane steroids substituted at the 3-, 6-, 19-, 20- and 21- positions. Many of the 3-substituted steroids are known compounds which may be derived from 3-oxo-steroids. As shown in Figure 1, prena-4,20-diene-3-one (1) can be converted to a 3, 5, 20-triene ether (2) or 1, 4, 20-trien-3-one (3), which are respective starting materials for 6- and 3- substituted hydroxy derivatives.

Alkoxy derivatives are prepared from their corresponding hydroxy steroids by reaction with an alkylating agent such as trimethyloxonium fluoroborate, triethyloxonium fluoroborate or methylfluorosulfonate in an inert chlorocarbon solvent such as methylene chloride. Alternatively, alkylating agents such as NaH, KM or KOBut, silver oxide or barium oxide in polar, aprotic solvents as for example, DMF, DMSO and hexamethylphosphoramide.

General procedures for synthetic reactions of steroids are known to those skilled in art. Where time and temperature of reactions must be determined, these can be determined by a routine methodology. After addition of the required reagents, the mixture is stirred under an inert atmosphere and aliquots are removed at hourly intervals. The aliquots are analyzed by chromatography to monitor the disappearance of starting material, at which point the work-up procedure is initiated. If the starting material is not consumed within twenty-four hours, the mixture is heated to reflux and hourly aliquots are analyzed, as before, until the starting material disappears. In this case the mixture

- 66 -

is allowed to cool before the work-up procedure is initiated.

Purification of the products is accomplished by means of chromatography and/or crystallization, as known to those skilled in the art.

2. Preparation of 19-OH derivatives.

Synthesis of 19-OH-pregna-4,17-diene-3-one.

A method of synthesizing this compound is provided in SCHEME 3.

10 D. Pharmaceutical Compositions and Methods of Use

An embodiment of the subject invention is a method of altering the hypothalamic function of an individual. Another embodiment is altering an autonomic function of an individual. These autonomic functions include but are not limited to heart rate, respiratory rate, brain wave patterns (percentage alpha cortical activity), body temperature. Other embodiments include, but are not limited to, methods of diminishing negative affect, negative mood or negative character traits of an individual. Another embodiment is a method of treating female premenstrual stress. All of these embodiments are accomplished by means of the non-systemic, nasal administration of certain pregnane or cholane steroids, combinations of pregnane and cholane steroids and combinations of one or more pregnane or cholane steroids and one or more androstane and/or estrene steroids.

This particular mode of administration is distinguished from alternative modes, such as ingestion or injection, in several important ways, these by virtue of the direct contact with the VNO provided by the nasal administration of the steroid ligand. In the methods of

- 67 -

this invention, the appropriate ligand is administered directly to the chemoreceptors in the nasal passage and the vomeronasal organ, without pills or needles - i.e., non-invasively. Drug action is mediated through binding of the ligands, described herein, to specific receptors displayed by neuroepithelial cells in the nose, preferably in the VNO. This furthermore, the mode of drug action is through the nervous system and not through the circulatory system - thus brain function can be affected without consideration of the blood-brain barrier. These methods of treatment provide a direct means of affecting the hypothalamus through the nervous system because there is only one synaptic junction between pheromone receptors and the hypothalamus. Because sensory nerves are addressed to a specific location in the brain, this method has a highly specific drug effect, thereby greatly reducing the potential of undesirable side-effects.

VNO contact is important because the VNO is associated with chemoreceptive/pheromonal function. The VNO consists of a pair of blind tubular diverticula which are found at the inferior margin of the nasal septum. The VNO contains neuro-epithelia, the axons of which have direct synapses to the amygdala and from there, to the hypothalamus. The existence of the VNO has been well documented in most terrestrial vertebrates including the human fetus; however, in adult humans it is generally thought to be rudimentary. (See Johnson, et al., supra.)

The ligand substances described herein, or their sulfated, cypionated, benzoated, proprionated, or glucuronated derivatives, may be administered directly, but are preferably administered as compositions. They are prepared in a liquid dosage form such as, for example, liquids, suspensions or the like, preferably in unit dosage forms suitable for single administration of

- 68 -

precise dosages. Liquid dosages may be administered as nose drops or as an aerosol. Alternatively, the active compound can be prepared as a creme or an ointment composition and applied topically within the nasal cavity. In addition, a vomeropherin may be administered as vapor contained in an air puff delivered to the nasal cavity. As another alternative, delivery may occur by controlled release of these agents by encapsulation either in bulk or at a microscopic level using synthetic polymers, such as silicone, and natural polymers such as gelatin and cellulose. The release rate can be controlled by proper choice of the polymeric system used to control the diffusion rate (Langer, R.S. and Peppas, N.A., Biomaterials 2,201, 1981). Natural polymers, such as gelatin and cellulose slowly dissolve in a matter of minutes to hours while silicone remains intact for a period of months. The compositions will include a conventional pharmaceutical carrier or excipient, one or more of the active pregnane or cholane compound(s) of Formula I, and the composition may or may not additionally include one or more androstane or estrene steroids. In addition, the compositions may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, etc.

The most likely means of communication of a semiochemical ligand is the inhalation of a naturally occurring pheromone present on the skin of another. Since these compounds are relatively nonvolatile, it is estimated that, even during intimate contact, a human subject would inhale picogram amounts of a naturally occurring steroid from the skin of another. From the amount inhaled it is estimated that only about 1% would reach the receptors of the vomeronasal organ.

The amount of vomeropherin administered will of course, be dependent on the subject being treated, the

- 69 -

severity of the affliction, the manner of administration, the frequency of administration, and the judgment of the prescribing physician. However, a single dosage of at least about 10 picograms, delivered directly into the lumen of the vomeronasal organ, is effective in eliciting a transient autonomic response. When administered to the nasal cavity, the dosage is about 100 picograms to about 100 micrograms, preferably about 1 nanogram to about 10 micrograms, more preferably about 10 nanograms to 1 about microgram. The frequency of administration is desirably in the range of an hourly dose to a monthly dose, preferably from 8 times/day to once every other day, more preferably 1 to 3 times per day. Ointments containing one or more active compounds and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like, can be prepared using a base such as, for example, petroleum jelly, lard, or lanolin.

Liquified pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, etc. an active compound as defined above and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, for example, sodium acetate, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, etc. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, PA, 15th Ed., 1975. The composition or formulation to be administered will, in

- 70 -

any event, contain a quantity of one or more of the active compound(s) in an amount effective to alleviate the symptoms of the subject being treated.

For aerosol administration, the active ingredient
5 is preferably supplied in finely divided form along with a surfactant and a propellant. Typical percentages of active ingredients are 0.001 to 2% by weight, preferably 0.004 to 0.10%.

Surfactants must, of course, be nontoxic, and
10 preferably soluble in the propellant. Representative of such agents are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, oleostearic and oleic acids with an aliphatic polyhydric
15 alcohol or its cyclic anhydride such as, for example, ethylene glycol, glycerol, erythritol, arabitol, mannitol, sorbitol, and hexitol anhydrides derived from sorbitol (the sorbitan esters sold under the trademark "Spans") and the polyoxyethylene and polyoxypropylene
20 derivatives of these esters. Mixed esters, such as mixed or natural glycerides, may be employed. The preferred surface-active agents are the oleates orbitan, e.g., those sold under the trademarks "Arlacel C" (sorbitan sesquioleate), "Span 80" (sorbitan monooleate) and "Span
25 85" (sorbitan trioleate). The surfactant may constitute 0.1-20% by weight of the composition, preferably 0.25-5%.

The balance of the composition is ordinarily propellant. Liquefied propellants are typically gases at ambient conditions, and are condensed under pressure.
30 Among suitable liquefied propellants are the lower alkanes containing up to five carbons, such as butane and propane; fluorinated or fluorochlorinated alkanes, such as are sold under the trademark "Freon". Mixtures of the above may also be employed.

- 71 -

In producing the aerosol, a container equipped with a suitable valve is filled with the appropriate propellant, containing the finely divided active ingredient and surfactant. The ingredients are thus maintained at an elevated pressure until released by action of the valve.

Yet another means of administration is topical application of a volatile liquid composition to the skin, preferably facial skin, of an individual. The composition will usually contain an alcohol such as ethanol or isopropanol. A pleasant odorant may also be included in the composition.

F. Measuring Affect, Mood and Character Trait.

Feeling states associated with affects, moods and character traits are generally measured by use of a questionnaire. For example questionnaires comprising a number of adjectives which refer to feeling states may be administered to an individual. The individual evaluates his or her feeling state described by the adjective and rates the intensity of the feeling on a numerical scale. Clustering of related adjectives and statistical analysis of a subject's evaluation of each adjective provides a basis for the measurement of various feeling states.

Alternatively, feeling states may be measured by autonomic changes, such as those used in polygraphic evaluations (galvanic skin response, pulse rate and the like). Cabanac, M. Annual Review of Physiology (1975) 37:415; Hardy, J.D., "Body Temperature Regulation", Chapter 59, pp. 1417. In: Medical Physiology. Vol. IIEd.: VB Mountcastle (1980); Wolfram Bouscein. Electrodermal Activity (Plenum Press 1992). In addition, non-verbal cues such as facial expression and body posture may be evaluated.


- 72 -

III. Examples


The following examples are intended to illustrate but not to limit the invention.


Abbreviations used in the examples are as follows:

s aq.=aqueous; RT.=room temperature; PE=petroleum ether
(b.p. 50-70°); DMF=N,N=dimethylformamide; DMSO=dimethyl
sulfoxide; THF=tetrahydrofuran.


SCHEME 1

- 74 -





SCHEME 2

SCHEME 1


SCHEME 4





Scheme 5 Oxidation of pregnanes




**Scheme 6 Synthesis of
21-methylene-20(R)-methylpregnanes**




Scheme 7 Synthesis of pregnanes.



Scheme 8 Synthesis of a cholatriene



Scheme 9 Synthesis of Cholenes



- 82 -

ExamplesExample 1 - Pregna-4, 20-dien-3 α -(β)-ol.

To a 1M solution of lithium trisiamylborohydride (5.0ml, 5.0mmole) at -78°C under argon was added a
5 solution of pregna-4,20-dien-3-one (1.10g, 3.70mmole) in dry THF (14ml), with stirring, and the mixture was allowed to warm to room temperature. After 3 hours, the mixture was cooled to -78°C and the following reagents were added sequentially: water (2ml), ethanol (6ml), 12%
10 aqueous KOH solution (10ml), and 3% hydrogen peroxide (50ml). The mixture was allowed to warm to room temperature with stirring. After 2 hours, ethyl acetate (200ml) was added and the stirring was continued. The organic layer was separated, and was washed with satd.
15 NaHSO₃ solution, satd. NaHCO₃ solution, and satd. NaCl solution, dried (Na₂SO₄), and evaporated in vacuo to give 2.1g crude material. This was purified by flash chromatography on 210g silica gel (230-400 mesh), eluting with EtOAc/CH₂Cl₂ (5:95-->7:93) to give three fractions.
20 Fraction 1 (0.8g) contained the impure 3 α -alcohol. Fraction 2 (0.1g) was a mixture of the 3 α - and 3 β -alcohols. Fraction 3 (0.25g) was the pure 3 β -alcohol (23%). Fraction 1 was repurified by flash chromatography on 80g silica gel (230-400 mesh), eluting with
25 EtOAc/hexane (10:90-->15:85) to give 0.15 g pure 3 α -alcohol (14%).

Example 2 - Pregna-3,5 20-trien-3-yl methyl ether, 2.

Referring to SCHEME 1, compounds 2, 3, 4 and 5 were prepared as follows.

30 A solution of pregna-4,20-dien-3-one (1, 1.00 g 3.35 mmol) in 2,2-dimethoxypropane (5.0 Ml, 41 mmol) dimethylformamide (5.0 mL) and methanol (0.2 mL) was refluxed with catalytic p-toluenesulfonic acid monohydrate (26.9 mg, 0.141 mmol) for 2 h. After

- 83 -

cooling, sodium bicarbonate (153.6 mg, 1.828 mmol) was added and the reaction mixture was partitioned between 75 mL of hexanes and 50 mL of ice water. The organic phase was washed twice with 50 mL portions of water and once
5 with 50 mL of brine, after which it was filtered through a 17 mm high x 30 mm diag. column of silica gel 60. Product was further eluted with 100 mL of hexanes. Concentration of the combined eluates and recrystallization from acetone/methanol gave lustrous
10 very slightly yellow platelets (828.7 mg, 2.652 mmol, 79%) m.p. 111-114°C.

Example 3 - Pregna-1,4,20-trien-3-one.

Pregna-4,20-dien-3-one (1, 1.19 g, 3.99 mmol) was refluxed for 24 h with 2,3-dichloro-5,6-dicyano-1,4-
15 benzoquinone (DDQ, 2.72 g, 12.1 mmol) in 40 mL of benzene under Argon. The cooled suspension was diluted with ether and washed with two 100 mL portions of 5% (w/w) sodium hydroxide, two 100 mL portions of water and once
20 with 100 mL of brine. Ether (100 mL) was added to the resulting emulsion, which was dried over sodium sulfate and then filtered through a column of sodium sulfate (20 g). After washing the residue twice with 50 mL portions of ether the combined filtrates were concentrated under
25 reduced pressure and then flash chromatographed (25% ethyl acetate/hexanes on silica gel) to give a slightly yellow crystalline solid (0.26 g, 0.88 mmol, 22%).

Example 4 - Pregna-1,4,20-trien-3-ol, 4.

Pregna-1,4,20-trien-3-one (3, 0.26 g, 0.88 mmol) in 25 mL of anh. ether was reduced under argon atmosphere
30 with lithium aluminum hydride (250.5 mg, 6.601 mmol) for 2 h and then quenched with 2.50 g of Glauber's salt. The resulting suspension was stirred 70 min., filtered, and washed twice with 50 mL portions of ether. After

- 84 -

concentrating the combined filtrates under reduced pressure the residue was purified using preparative TLC (35% ethyl acetate/hexanes on alumina to give white needles (26.1 mg, 87.4 μ mol, 10%) m.p. 98-128°C.

5 Example 5 - Pregna-4,20-dien-6 β -ol-3-one, 5.

m-Chloroperbenzoic acid (MCPBA, 77.4%, 763.4 mg, 3.42 mmol) suspended in 30 mL of 1,2-dimethoxyethane (DME), 6 mL of water and 2.4 mL of 5% (w/w) sodium hydroxide was added to a solution of pregna-3,5,20-trien-10 3-yl methyl ether (2, 400.3 mg, 1.281 mmol) in 20 mL of DME + 2 mL of water over 85 min. with stirring. The reaction was continued 5 h and was then poured into 50 mL of saturated sodium bicarbonate. The mixture was extracted three times with 50 mL of ether and the 15 combined organic extracts were washed with 50 g of 5% (w/w) sodium thiosulfate pentahydrate + three 50 mL portions of brine, dried over magnesium sulfate, and filtered through Celite. After washing the residue with 10 mL of ether the combined filtrates were concentrated 20 in vacuum. Flash chromatography (35% ethyl acetate/hexanes on silica gel) and preparative TLC (35% ethyl acetate/hexanes on silica gel) gave a difficultly separable mixture as white crystals (95.5 mg, 0.304 mmol, 24%).

25 Example 6 - 20,21 - Dimethylpregna-5,20-dien-3 β -ol, 6.

Referring to SCHEME 2, compounds 6, 7, 8 and 9 were prepared as follows.

Ethyltriphenylphosphonium bromide (25.99 g, 70.00 mmol) and potassium t-butoxide (7.86 g, 70.0 mmol) with 30 80 mL of anh. DMSO were stirred under Argon in oil bath at ca. 80°C for 1 h, after which pregn-5-en-3 β -ol-20-one (4.43 g, 14.0 mmol) in 80 mL of warm anh. DMSO was added. The red suspension was stirred 1 h, removed from

- 85 -

the heat and poured into 200 mL of ice-brine. The mixture was then extracted three times with 100 mL of ether and the combined organic extracts were washed with 100 mL of brine, dried over sodium sulfate, and filtered through Celite. After washing the residue with 50 mL of ether the combined filtrates were concentrated under reduced pressure. The yellow residue was taken up in 95% ethanol with heating, boiled briefly with 1 g of charcoal, and filtered through Celite. After cooling and filtration of the residue was recrystallized twice more from ethanol to give white crystals (1.746 g, 5.314 mmol, 38%), m.p. 140-145°C.

Example 7 - 20,21 - Dimethylpregna-4,20-dien-3,6-dione, 7.

Jones reagent (2.67 M, 2.0 mL, 5.3 mmol) was added to a solution of 20,21 -dimethylpregna-5,20-dien-3 β -ol (6, 460.1 mg, 1.400 mmol) in 50 mL of acetone and the reaction was stirred 45 min. After quenching with 2-propanol (1.0 mL) the mixture was poured into 100 mL of water and extracted three times with 50 mL of ethyl acetate. The combined organic extracts were washed with 50 mL of saturated sodium bicarbonate + 50 mL of brine, dried over magnesium sulfate, and filtered through Celite. The residue was washed with 25 mL of ethyl acetate and the combined filtrates were concentrated under reduced pressure. Flash chromatography (25% ethyl acetate/hexanes on silica gel) and recrystallization of the residue from 95% ethanol gave yellow needles (138.2 mg, 0.4059 mmol, 29%), m.p. 172-178°C.

Example 8 - 20,21 - Dimethylpregna-4,20-dien-3-one, 8.

20,21 -Dimethylpregna-5,20-dien-3 β -ol (6, 400.3 mg, 1.218 mmol) in 5 mL of methylene chloride was oxidized with pyridinium chlorochromate (525.4 mg 2.437

- 86 -

mmol) for 42 h. Ether (3.5 mL) was added and the suspension was filtered through a 5 mm dia. x 60 mm high column of silica gel. The column was further eluted with 3.5 mL of ether and the combined filtrates were concentrated under reduced pressure. Flash chromatography of the resin followed by recrystallization from aqueous ethanol gave yellow crystals (43.6 mg, 0.134 mmol 11%). m.p. 157-165°C.

Example 9 - Pregna-4,20-dien-3,6-dione 9.

10 A solution of pregna-5,20-dien-3 β -ol (300.5 mg, 1.000 mmol) in 35 mL of acetone was cooled in an ice water bath and 2.67 M Jones reagent (0.71 mL, 1.9 mmol) was added. After stirring 1 1/2 h a further 0.71 mL of Jones reagent were added and the reaction was continued
15 45 min, 2-Propanol (1.0 mL) was added and the mixture was poured into 100 mL of water. The mixture was then extracted twice with 50 mL of ethyl acetate and the combined organic extracts were washed with 50 mL of saturated sodium bicarbonate + 50 mL of water + 50 mL of
20 brine and filtered through a 21 mm dia x 22 mm high column of silica gel 60. The column was eluted further with 25 mL of ethyl acetate and the combined filtrates were concentrated under reduced pressure. Recrystallization of the residue from 95% ethanol gave a
25 light yellow powder (104.6 mg, 0.3348 mmol 33%), m.p. 114-120°C.

Example 10 - Pregna-5,17,20-trien-3 β -ol, 10.

A solution of 17 α -ethynylandrostenediol (439.4mg, 1.397 mmol) in 10mL of dry THF was added to a suspension
30 of lithium aluminum hydride (106.5 mg, 2.mmol) and aluminum chloride (122.9 mg, 0.9220 mmol) in 10 mL of dry THF under argon. After refluxing 17 h the reaction mixture was quenched by stirring 2 h with sodium sulfate

- 87 -

decahydrate (1.00 g, 3.10 mmol). The reaction was filtered and the residue washed with three 10 mL portions of THF. Concentration of the combined filtrates under reduced pressure gave 0.44 g of white solid, which was
5 purified by flash chromatography (30% ethyl acetate/hexanes on silica gel) and twofold recrystallization from aqueous ethanol, giving lustrous white crystals (92.0 mg, 0.303 mmol, 22%), m.p. 144-149°C.

10 Example 11 - 5 α -Chloro-6 β ,19-epoxypregn-17-en-3 β -ol, 11.

Referring to SCHEME 3, compounds 11, 12, 13, and 14 were made as follows:

Ethyltriphenylphosphonium bromide (3.05 g, 8.22 mmol) and potassium t-butoxide (0.92 g, 8.2 mmol) were
15 reacted under argon in anh. DMSO (9.2 mL) for 1 h in a 76-86°C bath, following which 5 α -chloro-6 β ,19-epoxyandrostan-3 β -ol-17-one (555.9 mg, 1.640 mmol) in 9.2 mL of warm anh. DMSO was added and the mixture stirred a further 1 h. The reaction was then poured into 25 mL of
20 ice-brine and extracted three times with 10 mL portions of ether. The combined organic extracts were washed with brine, dried over magnesium sulfate, and filtered through Celite. The residue was washed twice with 5 mL of ether and the combined filtrates were dried in vacuum. The
25 residual yellow oil was purified by flash chromatography (60% ethyl acetate/hexanes on silica gel) to give a white syrupy solid (0.34 g, 0.97 mmol, 59%).

Example 12 - 5 α -Chloro-6 β , 19-epoxypregn-17-en-3-one, 12.

A solution of 5 α -chloro-6 β ,19-epoxypregn-17-en-3 β -
30 ol (11, 0.34 g, 0.97 mmol) in 35 mL of acetone was cooled in an ice-acetone bath and 0.47 mL of 2.67M Jones reagent were added. After stirring 40 min. the reaction was quenched with the addition of 0.5 mL of 2-propanol.

- 88 -

Water (15 mL) was added, volatile components were removed under reduced pressure, and the mixture was extracted three times with 15 mL portions of methylene chloride. The combined organic extracts were washed with 15 mL of
5 saturated sodium bicarbonate + 15 mL of brine, dried over magnesium sulfate, and filtered through Celite. After washing the residue twice with 5 mL of methylene chloride the combined filtrates were dried in vacuum. The residue
10 was flash chromatographed on silica gel using 30% ethyl acetate/hexanes as eluent to give a white crystalline solid (0.34 g, 0.97 mmol, quantitative).

Example 13 - 6 β ,19-Epoxy pregn-4,17-dien-3-one, 13.

5 α -chloro-6 β ,19-epoxy pregn-17-en-3-one (12, 0.34 g, 0.97 mmol) was dissolved with warming in 10 mL of anh.
15 methanol, potassium acetate (0.60 g, 6.1 mmol) was added, and 6.5 mL of solvent were distilled off at room pressure. The residue was concentrated under reduced pressure, taken up in 25 mL of water, and extracted three times with 10 mL portions of methylene chloride. The
20 combined organic extracts were dried over magnesium sulfate and filtered through Celite. The residue was washed with 10 mL of methylene chloride and the combined filtrates concentrated under reduced pressure to give a white crystalline solid (290.0 mg, 0.9281 mmol, 96%)
25 homogeneous to TLC (60% ethyl acetate/hexanes on silica gel; R_f 0.61).

Example 14 - Pregna-4,17-dien-19-ol-3-one, 14.

To a solution of 6 β ,19-epoxy pregna-4,17-dien-3-one (290.0 mg, 0.9281 mmol) in 10 mL of glacial acetic acid
30 was added zinc dust (1.12 g, 17.1 mg-atom) activated by stirring 2 min. with 10% hydrochloric acid followed by washing with water and acetone. The suspension was stirred vigorously for 10 min. in a 99-102°C and was then

- 89 -

filtered through Celite. The residue was washed 4 times with 10 mL of acetic acid and the combined filtrates were concentrated in vacuum. The residue was taken up in 50 mL of ethyl acetate, washed with 50 mL of water + 50 mL of saturated sodium bicarbonate + 50 mL of brine, dried over magnesium sulfate, and filtered through Celite. The residue was washed with 10 mL of ethyl acetate and the combined filtrates dried in vacuum. The residue was recrystallized from ethyl acetate to give white crystals (46.4 mg, 0.148 mmol, 16%), m.p. 192-195°C.

Example 15 - Pregn-4-en-3 β -ol-20-yne, 15.

Pregn-4-en-3 β -ol-20-yne, 15: Pregna-4-en-3-on-20-yne (200.1 mg, 0.6750 mmol) and lithium aluminum tri(t-butoxy)hydride (343.8 mg, 1.352 mmol) were suspended in 3.6 mL of anhydrous ether. After reacting 4 h, a further 343.5 mg (1.351 mmol) of hydride were added and the reaction was allowed to continue 16 h. See Scheme 4. After quenching with sodium sulfate decahydrate (3.41 g) the reaction mixture was agitated 15 min. and then filtered through diatomaceous earth. The residue was extracted 5 times with 10 mL portions of ether and the combined filtrates were concentrated under reduced pressure. Flash chromatography of the residue (25% ethyl acetate/hexanes on silica gel) followed by recrystallization from aqueous ethanol yielded a white powder (85.0 mg, 0.285 mmol, 42%), m.p. 120.5 - 123.5°C.

Example 16 - 20,21-Dimethylpregna-4,20E-diene-3,6-dione, 16.

Jones reagent (2.67 M, 1.75 mL, 4.67 mmol) was added to a solution of 20,21-dimethylpregna-5,20E-dien-3 β -ol (400.0 mg, 1.281 mmol) in 45 mL of acetone and the reaction was stirred 30 min. The mixture was then poured into 85 mL of water and extracted three times

- 90 -

with 40 mL portions of ethyl acetate. The combined organic extracts were washed with 40 mL of saturated sodium bicarbonate + 40 mL of brine, dried over magnesium sulfate, and filtered through diatomaceous earth. The residue was washed with 10 mL of ethyl acetate and the combined filtrates were concentrated under reduced pressure. Flash chromatography (20% ethyl acetate/hexanes on silica gel) of the residue followed by recrystallization from aqueous ethanol gave light yellow needles (119.1 mg, 0.3497 mmol, 29%), m.p. 171-173°C. TLC (20% ethyl acetate/hexanes on silica gel) showed a major component at R_f 0.17 with a minor contaminant at R_f 0.24.

Example 17 - 20-Methylpregna-4,20-diene-3,6-dione, 17.

Jones reagent (2.67 M, 1.83 mL, 4.89 mmol) was added to a solution of 20-methylpregna-5,20-dien-3 β -ol (400.0 mg, 1.272 mmol) in 45 mL of acetone and the reaction was stirred 20 min. After quenching with 2-propanol (0.91 mL) the mixture was poured into 85 mL of water and extracted three times with 40 mL portions of ethyl acetate. The combined organic extracts were washed with 40 mL of saturated sodium bicarbonate + 40 mL of brine, dried over magnesium sulfate, and filtered through diatomaceous earth. The residue was washed with 25 mL of ethyl acetate and the combined filtrates were concentrated under reduced pressure. Recrystallization of the solid residue from aqueous ethanol gave yellow crystals (235.0 mg, 0.7298 mmol, 57%), m.p. 148-150°C. TLC (20% ethyl acetate/hexanes on silica gel) showed a major product at R_f 0.24 with a minor impurity at the origin.

- 91 -

Example 18 - Pregna-4,16-diene-3,6-dione, 18.

Jones reagent (2.67 M, 0.23 mL, 0.61 mmol) was added to a solution of pregna-5,16-dien-3 β -ol (45.5 mg, 0.151 mmol) in 5 mL of acetone and the mixture was stirred 20 min. After quenching with 2-propanol (0.11 mL) 10 mL of water were added and the mixture was extracted three times with 5 mL portions of ethyl acetate. The combined organic extracts were washed with 5 mL of saturated sodium + 5 mL of brine, dried over magnesium sulfate, and filtered through diatomaceous earth. The residue was washed with 5 mL of ethyl acetate and the combined filtrates were concentrated under reduced pressure. Preparative TLC (25% ethyl acetate/hexanes on silica gel) of the residue gave a light yellow solid (10.6 mg, 33.9 μ mol, 22%) homogeneous to TLC (R_f 0.41, 25% ethyl acetate/hexanes on silica gel; starting material R_f 0.30).

Example 19 - Pregna-4,17,20-triene-3,6-dione, 19.

Jones reagent (2.67 M, 1.03 mL 2.75 mmol) was added to pregna-5,17,20-trien-3 β -ol (214.5 mg, 0.7187 mmol) in 25 mL of acetone and the mixture was stirred 20 min. After quenching with 2-propanol (0.52 mL), the mixture was poured into 50 mL of water and extracted three times with 25 mL portions of ethyl acetate. The combined organic extracts were washed with 25 mL of saturated sodium bicarbonate + 25 mL of brine, dried over magnesium sulfate, and filtered through diatomaceous earth. The residue was washed with 10 mL of ethyl acetate and the combined filtrates were concentrated under reduced pressure. Recrystallization with concomitant treatment with charcoal in aqueous ethanol gave light yellow needles (67.7 mg, 0.218 mmol, 30%), m.p. 140-143. TLC (25% ethyl acetate/hexanes on silica gel) shows product at R_f 0.43 with minor contaminants at R_f

- 92 -

0.86, 0.14, 0.08, and 0.00 (pregna-5,17-dien-3 β -ol R_f 0.32).

Example 20 - 21-Methylene-20(R)-methylpregn-4-en-3-one, 20.

5 Aluminum isopropoxide (0.39 g, 1.9 mmol) in 3.6 mL of hot toluene was added to a solution of 21-methylene-20(R)-methylpregn-5-en-3 β -ol (208.2 mg, 0.6337 mmol) in cyclohexanone (3.6 mL, 35 mmol) and 18 mL of toluene, and the mixture was refluxed with exclusion of moisture for 2
10 h. After cooling in ice, 0.92 mL of water and 2.2 mL of 3.6 N sulfuric acid were added and the mixture was shaken 1 min. Further water (4.6 mL) was added, the mixture was shaken 5 min., and the aqueous phase was discarded. Volatile components were removed by azeotropic/steam
15 distillation and the resulting aqueous suspension was extracted three times with 5 mL portions of methylene chloride. The combined organic extracts were filtered through a small column of sodium sulfate resting on a bed of diatomaceous earth, all of which was contained in a
20 Pasteur pipette. The residue was washed with 5 mL of methylene chloride and the combined filtrates were concentrated under reduced pressure. Recrystallization of the residual film gave white needles (152.7 mg, 0.4677 mmol, 74%), m.p. 138-139°C. TLC (25% ethyl
25 acetate/hexanes on silica gel) showed a major product at R_f 0.54 with a trace contaminant at R_f 0.71 (starting material R_f 0.36).

Example 21 - 21-Methylene-20(R)-methylpregn-4-en-3 β -ol, 21.

30 A suspension of 21-methylene-20(R)-methylpregn-4-en-3-one (5, 100.0 mg, 0.3063 mmol) and lithium tri-*t*-butoxyaluminumhydride (319.4 mg, 1.256 mmol) in 5 mL of anhydrous THF was stirred under argon for 6 h, after which

- 93 -

water (48 μ L), 15% (w/w) sodium hydroxide (48 μ L), and water (143 μ L) were added. The mixture was filtered through diatomaceous earth and the residue was extracted four times with 5 mL aliquots of THF. Concentration of the combined filtrates under reduced pressure and two-fold recrystallization from aqueous ethanol gave white platelets (42.6 mg, 0.130 mmol, 42%), m.p. 121-123°C. TLC (25% ethyl acetate/hexanes on silica gel) shows major product at R_f 0.38 and minor product at R_f 0.43 (starting material R_f 0.50).

Example 22 - 21-Methylene-20(R)-methylpregn-4-ene-3,6-dione, 22.

Jones reagent (2.67 M, 0.68 mL, 1.8 mmol) was added to 21-methylene-20(R)-methylpregn-5-en-3 β -ol (149.8 mg, 0.4560 mmol) in 15 mL of acetone and the reaction was stirred 20 min. After quenching with 2-propanol (0.34 mL), the mixture was poured into 30 mL of water and extracted three times with 15 mL portions of ethyl acetate. The combined organic extracts were washed with 15 mL of saturated sodium bicarbonate + 15 mL of brine, dried over magnesium sulfate, and filtered through diatomaceous earth. The residue was washed with 10 mL of ethyl acetate and the combined filtrates were concentrated under reduced pressure. Preparative TLC of the residue (25% ethyl acetate/hexanes on silica gel GF, 1000 μ) gave a yellow, crystalline solid (49.7 mg, 0.146 mmol, 32%) homogeneous to TLC (25% ethyl acetate/hexanes; R_f 0.47; starting material R_f 0.35).

Example 23 - Pregn-4-en-3 β -ol-20-yne, 23.

Lithium tri-*t*-butoxyaluminumhydride (343.8 mg, 1.352 mmol) was added to pregn-4-en-3-on-20-yne (200.1 mg, 0.6750 mmol) suspended in 3.6 mL of anhyd. ether and the mixture was stirred 4 h. Additional hydride (343.5

- 94 -

mg, 1.351 mmol) was added and the reaction was stirred a further 16 h. Glauber's salt (3.41 g) was added and the suspension was agitated 15 min. The mixture was filtered through diatomaceous earth and the residue was extracted 5 five times with 10 mL portions of ether. Concentration of the combined filtrates under reduced pressure followed by flash chromatography (25% ethyl acetate/hexanes on silical gel) and recrystallization from aqueous ethanol gave a white powder (85.0 mg, 0.285 mmol, 42%), m.p. 10 120.5-123.5°C. TLC (25% ethyl acetate/hexanes on silica gel) showed product (R_f 0.23) contaminated with traces of what appeared to be starting material (R_f 0.29).

Example 24 - Pregn-4,16-dien-6 β -ol-3-on-20-yne, 24.

3-chloroperbenzoic acid (290.0 mg, 1.680 mmol) in 15 9.4 mL of 1,2-dimethoxyethane + 3.6 mL of water was added to a suspension of pregna-3,5,16-trien-20-yn-3-yl methyl ether (471.0 mg, 1.527 mmol) in 1,2-dimethoxyethane (9.4 mL) and the reaction was stirred 30 min. The mixture was poured into 50 mL of saturated sodium bicarbonate and 20 extracted three times with 50 mL of ethyl acetate. The combined organic extracts were washed with 50 g of 5% (w/w) sodium thiosulfate pentahydrate + three 50 mL aliquots of brine, dried over magnesium sulfate, and filtered through diatomaceous earth. The residue was 25 washed with 25 mL of ethyl acetate and the combined filtrates were concentrated under reduced pressure. Flash chromatography (50% ethyl acetate/hexanes on silica gel) of the residue, followed by recrystallization from aqueous ethanol gave a light yellow solid (145.9 mg, 30 0.4700 mmol, 31%), m.p. >300°C.

Example 25 - Chola-5,16,20(22)-trien-3 β -ol, 25.

A suspension of propyltriphenylphosphonium bromide (12.13 g, 31.48 mmol) and potassium t-butoxide (3.54 g,

- 95 -

31.5 mmol) in anh. DMSO (35 mL) under argon atmosphere was placed in an oil bath (72-87°C) and then stirred 1 h. Pregna-5,16-dien-3 β -ol-20-one (1.9807 g, 6.298 mmol) in 35 mL of warm anh. DMSO was added and the reaction was stirred 90 min. The mixture was then poured into 90 mL of ice-brine and extracted with three 45 portions of ether. The combined organic extracts were washed with 45 mL of brine, dried over sodium sulfate, and filtered. The residue was washed with 10 mL of ether and the combined filtrates were concentrated under reduced pressure. Flash chromatography (20% ethyl acetate/hexanes on silica gel) of the resulting oil gave an amber resin (103.6 mg, 0.3042 mmol, 4.8%).

Example 26 - Chola-4-20(22)E-diene-4,6-dione, 26.

Jones reagent (2.67 M, 1.26 mL, 3.36 mmol) was added to chola-5,20(22)E-dien-3 β -ol (300.0 mg, 0.8758 mmol) in 30 mL of acetone and the reaction was stirred 20 min. After quenching with 2-propanol (0.63 mL), the mixture was poured into 60 mL of water and extracted three times with 30 mL portions of ethyl acetate. The combined organic extracts were washed with 30 mL of saturated sodium bicarbonate + 30 mL of brine, dried over magnesium sulfate, and filtered through diatomaceous earth. The residue was washed with 10 mL of ethyl acetate and the combined filtrates were concentrated under reduced pressure. Recrystallization from aqueous ethanol gave a yellow solid (139.3 mg, 0.3929 mmol, 45%), m.p. 109-116°C. TLC (25% ethyl acetate/hexanes on silica gel) shows a major product (R_f 0.45) with contaminants at R_f 0.24, 0.1, and 0.08 (chola-4,20(22)E-dien-3-one R_f 0.49).

- 96 -

Example 27 - Chola-4,20(22)E-dien-3-one, 27.

Aluminum isopropoxide (2.82 g, 13.8 mmol) in 26 mL of hot toluene was added to chola-5,20(22)E-dien-3 β -ol (1.5777 g, 4.605 mmol) in cyclohexanone (26 mL, 0.25 mol) + 130 mL of toluene and the reaction was refluxed 4 h with exclusion of moisture. After cooling, water (6.7 mL) and 3.6 N sulfuric acid (16 mL) were added and the mixture was shaken 1 min. Further water (32 mL) was added, the mixture was shaken 5 min., and the aqueous layer was discarded. The volatile components were removed by azeotropic/steam distillation and the resulting aqueous suspension was extracted three times with 20 mL portions of methylene chloride. The combined organic extracts were dried over magnesium sulfate and filtered through diatomaceous earth. The residue was washed with 10 mL of methylene chloride and the combined filtrates were concentrated under reduced pressure. Recrystallization from aqueous ethanol with concomitant treatment with charcoal gave light yellow crystals (1.3317 g, 3.9104 mmol, 85%), m.p. 133-135°C. TLC (25% ethyl acetate/hexanes on silica gel) showed the product was homogeneous (R_f 0.49; starting material R_f 0.31).

Example 28 - Chola-4,20(22)E-dien-3 β -ol, 28.

A suspension of chola-4,20(22)E-dien-3-one (3, 400.0 mg, 1.175 mmol) and lithium tri-*t*-butoxyaluminumhydride (1.2250 g, 4.8158 mmol) in 15 mL of anhydrous ether was stirred under argon for 8 h. Glauber's salt (6.07 g) was added, the mixture was stirred 5 min., and was then filtered through diatomaceous earth. The residue was extracted five times with 15 mL portions of ether and the combined filtrates were concentrated under reduced pressure. Preparative TLC (5% ethyl acetate/methylene chloride on silica gel, 1000 μ) of the residue gave a light yellow solid (49.9 mg, 0.146 mmol,

- 97 -

12%) homogeneous to TLC (5% ethyl acetate/methylene chloride on silica gel, R_f 0.34; pregna-5,20-dien-3 β -ol R_f 0.26).

Example 29 - Chola-3,5,20(22)E-trien-3-yl methyl ether.

5 29.

A mixture of chola-4,20(22)E-dien-3-one (808.6 mg, 2.374 mmol), 2,2-dimethoxypropane (3.1 mL, 25 mmol), dimethylformamide (3.1 mL), methanol (0.13 mL), and p-toluenesulfonic acid monohydrate (28.0 mg, 0.147 mmol) was refluxed 4 h with exclusion of moisture. After cooling, sodium bicarbonate (0.17 g) was added and the mixture was partitioned between 30 mL of water and 50 mL of ether. The organic layer was washed with 30 mL of brine, dried over magnesium sulfate, and filtered through diatomaceous earth. The residue was washed with 10 mL of ether and the combined filtrates were concentrated under reduced pressure. Recrystallization of the resulting solid from acetone gave a light yellow solid (453.8 mg, 1.280 mmol, 54%), m.p. 94-96°C. TLC (10% ethyl acetate/hexanes on silica gel) showed product at R_f 0.62 with a trace of starting material (R_f 0.14).

Example 30 - Chola-4,20(22)E-dien-6 β -ol-3-one, 30.

3-Chloroperbenzoic acid (183.0 mg, 1.060 mmol) in 5.9 mL of 1,2-dimethoxyethane + 2.3 mL of water was added to chola-3,5,20(22)E-trien-3-yl methyl ether in 5.9 mL of 1,2-dimethoxyethane over 2 1/2 min. After stirring the reaction a further 1/2 h, it was poured into 30 mL of saturated sodium bicarbonate and extracted three times with 30 mL portions of ethyl acetate. The combined organic extracts were washed with 30 g of 5% (w/w) sodium thiosulfate pentahydrate + three 30 mL portions of brine, dried over magnesium sulfate, and filtered through diatomaceous earth. The residue was washed with 10 mL of

- 98 -

ethyl acetate and the combined filtrates were concentrated under reduced pressure. Flash chromatography (60% ethyl acetate/hexanes on silica gel) of the resulting solid and subsequent preparative TLC (50% ethyl acetate/hexanes on silica gel) of the
5 resulting solid and subsequent preparative TLC (50% ethyl acetate/hexanes on silica gel, 1000 μ) gave a white crystalline solid (114.3 mg, 0.3206 mmol, 33%) consisting of two components by TLC (50% ethyl acetate/hexanes on
10 silica gel; major R_f 0.48, minor R_f 0.41; chola-4,20(22)E-dien-3-one R_f 0.74).

Example 31 - Measurement of Autonomic Responses to Stimulation of the VNO.

Various autonomic parameters were monitored as

- 15 A1-P3 Pregna-4,16-dien-3-one
- A2-P3 Pregna-5,16-dien-3 β -ol
- A8-P1 3-Methoxy-pregna-3,5,20-triens
- A6-P1 Pregna-4,20-dien-3,6-dione
- 20 20,21-Dimethylpregna-5,20-dien-3 β -ol
- 20,21-Dimethylpregna-5,20-dien-3-one
- A14-P2 6 β ,19-Epoxypregna-4,17-dien-3-one
- A7-P2 19-Hydroxy-pregna-4,17(20)-dien-3-one
- A13-P1 Pregna-4,20-dien-6 β -ol-3-one
- A11-P1 Pregna-1,4,20-trien-3-one
- 25 A1-P1 Pregna-4,20-dien-3-one
- A2-P1 Pregna-5,20-dien-3 β -ol
- A4-P1 Pregna-4,20-3 α -ol
- A3-P1 Pregna-4,20-3 β -ol
- A1-P4 Pregn-4-en-3-one
- 30 A2-P4 Pregn-5-en-3 β -ol

was administered to 24 female and 24 male subjects using the procedure described in Example 32. Propylene glycol was also administered as a control. When compared to a propylene glycol control, the test compounds induced a

- 99 -

significant change in the integrated receptor potential in the VNO, galvanic skin response (GSR), skin temperature (ST), the percentage of cortical alpha wave activity as measured by electroencephalogram (EEG), electrocardiogram (EKG) and respiratory frequency (RF). The results are shown in FIGS. 2 through 58. Additional tests were performed on other pregnane derivatives, the results of which are shown in FIGS. 59-84. The results of tests on cholane derivatives are shown in FIGS. 85-96.

10 Example 32 - Electrophysiological Studies

The following electrophysiological studies were performed in clinically normal human volunteers of both sexes whose ages ranged from 20 to 45 years. No anesthetics were used, and female subjects were excluded if pregnant.

The stimulation and recording system consists of a "multifunctional miniprobe" described elsewhere (Monti-Bloch, L. and Grosser, B.1. (1991) "Effect of putative pheromones on the electrical activity of the human vomeronasal organ and olfactory epithelium," J. Steroid Biochem. Molec. Biol. 39:573-582.). The recording electrode is a 0.3 mm silver ball attached to a small (0.1 mm) silver wire insulated with Teflon® the surface of the electrode is first treated to produce a silver chloride interface, and is then covered with gelatin. It is positioned within a small caliber Teflon® catheter (dia = 5 mm) such that the tip of the electrode protrudes approximately 2 mm. The Teflon® catheter is 10 cm in length and constitutes the terminal extension for a multichannel delivery system which delivers a continuous air stream carrying discreet pulses of chemosensory stimuli. The air stream first passes into a small chamber and is bubbled through a solution containing either a vomeropherin or an olfactant in a diluent or the

- 100 -

diluent alone. A solenoid is used to rapidly redirect the air stream from the chamber to a route which bypasses the chamber. This creates a discreet pulse of stimulant in the air stream. A second, outer Teflon® tube with a diameter of 2 mm surrounds the catheter-electrode assemblage, and its central end is connected to an aspirator that provides continuous suction of 3ml/s. This concentric arrangement of the outer suction tube allows the emitted chemosensory stimuli to be localized to an area we call a "minifield" (approx. dia = 1 mm), and it avoids diffusion of substances either to the area outside the intended stimulation site or into the respiratory system. The entire stimulating and recording assemblage may be positioned either on the neurosensory epithelium within the VNO, or on the surface of the olfactory or respiratory epithelium.

Electro-vomeronasogram (EVG): Recordings are carried out in a quiet room with the subject supine; the multi-functional miniprobe is initially stabilized within the nasal cavity using a nasal retractor placed in the vestibule. Reference and ground electrodes consist of silver discs (8 mm), both of which are positioned on the glabella.

The entrance to the VNO, or vomeronasal pit, is identified by first dilating the nasal aperture and vestibule. A 6x magnifying binocular loupe with halogen illumination is then used to introduce the tip of the Teflon® catheter and recording electrode assemblage into the VNO opening where it is stabilized at an approximate depth of 1 mm within the vomeronasal passage. Optimal placement of the recording electrode is signaled after testing for an adequate depolarization in response to a test substance.

- 101 -

Electrical signals from the recording electrode are fed to a DC amplifier after which they are digitized, computer monitored, and stored. The peak-to-peak amplitude of the signals is measured, and the area under the depolarization wave is integrated, while continuously monitoring the signal both on the computer screen and on a digital oscilloscope. The integrated EVG is shown in FIGs 1 and 2 for compounds A1-P1, A2-P1, A4-P1, A3-P1, A1-P4, A2-P4 (referring to the chart). Artifacts produced by respiratory movements are deleted by training the subjects to practice mouth breathing with velopharyngeal closure. Samples of vomeropherins in concentration of 25-800 fmoles are delivered in the continuous air stream for durations from 300 milliseconds to 1 second. Usually, intervals of 3 to 5 minutes separated each series of short test pulses. All components of the lines carrying the test stimuli are made of Teflon®, glass or stainless steel and are carefully cleaned and sterilized before each use. Activity was recorded using standard electroencephalographic (EEG) electrodes placed at positions Cz-A1 and Tz-A1 of the international 10120 system; the ground electrode was placed on the mastoid process. Skin temperature (ST) was recorded by a small (1.0 mm) thermistor probe placed in the right ear lobe. Respiratory frequency (RF) was measured with an adjustable strain gauge placed around the lower thorax. All electrical signals were DC amplified, digitized (MP-100, Biopac Systems) and continuously monitored utilizing a computer.

Statistical Analysis: EVGs, peak-to-peak changes and frequency changes of other parameters were measured and statistically analyzed. The significance of the

- 102 -

results was determined by either using paired t-tests or analysis of variance (ANOVA)..

Reflex Effects of Vomeropherins: Studies were conducted to determine the central nervous system (CNS) reflex responses to vomeropherin stimulation of the VNO. The sexually dimorphic local responses induced by vomeropherins were mirrored in the autonomic response of male and female subjects.


Cortical activity was recorded from Cz and Tz in male and female subjects during application to the VNO of air pulses (300 ms to 1 sec) containing 200 fmoles of vomeropherin. There is also preliminary evidence that the EVG is not associated with trigeminal nociceptor endings since application of a local anesthetic (2% lidocaine) to the respiratory epithelium of the nasal septum neither blocks nor diminishes the EVG (Monti-Bloch, L. and Grosser, B.I. (1991) "Effect of putative peromones on the electrical activity of the human vomeronasal organ and olfactory epithelium," J. Steroid Biochem. Molec. Biol. 39:573-582), also subjects failed to report sensations of pain as a consequence of any of the stimulation procedures.

- 103 -

CLAIMS

What is claimed:

1. A pharmaceutical composition suitable for nasal administration in an individual, said composition comprising a steroid and a pharmaceutically acceptable carrier, wherein said steroid has the formula:



- wherein P₁ is selected from the group consisting of oxo, α-(β-) hydroxy, α-(β-) acetoxy, α-(β-) propionyloxy, α-(β-) methoxy, α-(β-) lower acyloxy, α-(β-) lower alkyloxy, and α-(β-) benzyloxy; P₂ is selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl, alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxyalkyl; P₃ is selected from the group consisting of hydrogen, oxo, halo, hydroxy, alkoxy, and acyloxy; P₄ through P₁₂ may each be, independently, hydrogen, halo, methyl, or halo-, dihalo- or perhalomethyl; P₁₃ is hydrogen, methyl, methylene, halo-substituted methyl or halo-substituted methylene, ethyl, ethylenyl, acetylenyl, methyl-methylenyl or methyl-methinylenyl; and "a", "b", "c", "d", "e", "h", "i", "j" and "k" are alternative sites for optional double bonds and "j" or "k" may also be triple bonds; and when P₂ is methyl and P₃ is β-hydroxy, P₂ and P₃ may be joined to form a cyclic ether.

- 104 -

2. A composition according to Claim 1 wherein "b" is a double bond.

3. A composition according to Claim 2 wherein "e" or "d" is a double bond.

5 4. A composition according to Claim 1 wherein "a" and "c" are double bonds.

5. A composition according to Claim 1 wherein "h" is an optional double bond, and "i" and "j" are absent.

10 6. A composition according to Claim 1 wherein "j" is a double bond.

7. A composition according to Claim 1 wherein "j" is a triple bond.

8. The composition according to Claim 1 wherein
15 P_{13} is hydrogen, methyl, methylene, halo-substituted methyl or halo-substituted methylene.

9. The composition according to Claim 1 wherein P_{13} is ethyl, ethylenyl, acetylenyl, methyl-methylenyl, or methyl-methinyl.

20 10. The pharmaceutical composition of any of Claims 1 through 9 wherein said steroid is dissolved in said carrier.

11. The pharmaceutical composition of any of Claims 1 through 9 wherein said composition is in a
25 liquid form.

- 105 -

12. The pharmaceutical composition of any of Claims 1 through 9 wherein said composition further contains a pharmaceutically acceptable ointment base.

13. The pharmaceutical composition of any of Claims 1 through 9 which contains no more than one of said steroids.

14. The pharmaceutical composition of any of Claims 1 through 9 which contains more than one of said steroids.

15. A method of altering a hypothalamic function of an individual, said method comprising:

providing a steroid selected from the group consisting of pregnanes, cholanes and their derivatives on the surface of nasal neuroepithelial cell of said individual wherein said cell is a part of tissue other than olfactory, epithelia; and,

administering said steroid within a nasal passage of said individual such that said pregnane derivative binds specifically to said receptor and results in an alteration of hypothalamic function of said individual.

16. A method of altering an autonomic function of an individual, said method comprising:

providing a steroid selected from the group consisting of pregnanes, cholanes and their derivatives for a chemoreceptor of a nasal neuroepithelial cell of said individual


- 106 -

wherein said cell is a part of tissue other than olfactory epithelia; and,

administering said steroid within a nasal passage of said individual such that said
 5 ligand binds specifically to said receptor and results in an alteration of hypothalamic function of said individual.

17. The method of Claim 15 or 16 wherein said neuroepithelial cell is located within a vomeronasal
 10 organ of said individual.

18. The method of Claim 17 wherein said pregnane is of the formula:



wherein P₁ is selected from the group consisting of oxo,
 α-(β-) hydroxy, α-(β-) acetoxy, α-(β-) propionyloxy,
 15 α-(β-) methoxy, α-(β-) lower acyloxy, α-(β-) lower
 alkyloxy, and α-(β-) benzoyloxy; P₂ is selected from the
 group consisting of methyl, hydroxymethyl, acyloxymethyl,
 alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl,
 and alkoxylalkyl; P₃ is selected from the group consisting
 20 of hydrogen, oxo, halo, hydroxy, alkoxy, and acyloxy; P₄
 through P₁₂ may each be, independently, hydrogen, halo,

- 107 -

methyl, or halo-, dihalo- or perhalomethyl; P_3 is hydrogen, methyl, methylene, halo-substituted methyl or halo-substituted methylene, ethyl, ethylenyl, acetylenyl, methyl-methylenyl or methyl-methinyl; and "a", "b", "c",
5 "d", "e", "h", "i", "j" and "k" are alternative sites for optional double bonds and "j" or "k" may also be triple bonds; and when P_2 is methyl and P_3 is β -hydroxy, P_2 and P_3 may be joined to form a cyclic ether.

19. A method according to Claim 18 wherein "b" is
10 a double bond.

20. A method according to Claim 19 wherein "e" or "d" is a double bond.

21. A method according to Claims 18 wherein "a" and "c" are double bonds.

15 22. A method according to Claim 18 wherein "h" is an optional double bond, and "i" and "j" are absent.

23. A method according to Claim 18 wherein is a double "j" is a double bond.

24. A method according to Claim 18 wherein "j" is
20 a triple bond.

25. The method of Claim 17 wherein the amount of said pregnane derivative that is administered is at least about 100 picograms, but no more than about 100 micrograms.

25 26. The method of Claim 25 wherein the amount of said pregnane derivative that is administered is at least about 1 nanograms, but no more than about 10 micrograms.

- 108 -

27. The method of Claim 26 wherein the amount of said pregnane derivative that is administered is at least about 10 nanograms, but no more than about 1 microgram.

28. The method of Claim 15 or 16 further
s comprising one step of preparing a pharmaceutical composition of said pregnane derivative dissolved in a pharmaceutically acceptable carrier.

29. The method of Claim 28 wherein said pharmaceutical composition is an ointment.


10 30. The method of Claim 28 wherein said pharmaceutical composition is liquid.

31. The method of Claim 28 wherein the administration is by aerosol.

32. The method of Claim 15 or 16 wherein more
15 than one steroid is administered.

- 109 -

33. A compound of the formula



wherein P₁ is selected from the group consisting of oxo, α-(β-) hydroxy, α-(β-) acetoxy, α-(β-) propionyloxy, α-(β-) methoxy, α-(β-) lower acyloxy, α-(β-) lower alkyloxy, and α-(β-) benzoyloxy; P₂ is selected from the group consisting of methyl, hydroxymethyl, acyloxymethyl, alkoxymethyl, lower alkyl, hydroxyalkyl, acyloxyalkyl, and alkoxyalkyl; P₃ is selected from the group consisting of hydrogen, oxo, halo, hydroxy, alkoxy, and acyloxy; P₄ through P₁₂ may each be, independently, hydrogen, halo, methyl, or halo-, dihalo- or perhalomethyl; P₁₃ is hydrogen, methyl, methylene, halo-substituted methyl or halo-substituted methylene, ethyl, ethylenyl, acetylenyl, methyl-methylenyl or methyl-methinyl; and "a", "b", "c", "d", "e", "h", "i", "j" and "k" are alternative sites for optional double bonds and "j" or "k" may also be triple bonds; and when P₂ is methyl and P₃ is β-hydroxy, P₂ and P₃ may be joined to form a cyclic ether; with the provisos that

- i) when P₁ = oxo; b is present; e, a, c, d are absent; P₂ = methyl; and P₃, P₄, P₇, P₉, P₁₀, P₁₁ and P₁₂ are hydrogen; then if h is present or absent, P₆ must be present and cannot be hydrogen;

- 110 -

- ii) when $P_1 = \text{OH}$ and c is present; e , a , b and d are absent; $P_2 = \text{methyl}$; and
 P_3 , P_4 , P_7 , P_9 , P_{10} , P_{11} , P_{12} and P_{13} are hydrogen; then
- 5 (a) if h is present, P_6 must be present and is not hydrogen;
(b) if h is absent and P_6 is present, P_6 cannot be hydrogen;
(c) if h is absent and P_6 is present,
10 then i and j are both present;
- iii) when $P_1 = \beta\text{-OH}$ and b is present; e , a , c , d are absent; $P_2 = \text{methyl}$; and
 P_3 , P_4 , P_7 , P_9 , P_{10} , P_{11} and P_{13} are hydrogen;
- 15 (a) if h is present, i and/or j must be present; or P_6 is not hydrogen;
(b) if h is present, j cannot be a triple bond;
- iv) when $P_1 = \alpha\text{-OH}$; b is present; e , a , c , d are absent; $P_2 = \text{methyl}$;
20 and P_3 , P_4 , P_7 , P_9 , P_{10} , P_{11} and P_{13} are hydrogen; then
(a) if h is present, i must be absent and P_6 cannot be hydrogen;
25 (b) if i and j are both absent, P_6 cannot be hydrogen;
- v) when $P_1 = \text{oxo}$; and b and d are present; e , a , and c are absent; $P_2 = \text{methyl}$;
and P_3 , P_4 , P_7 , P_9 , P_{11} and P_{13} are all
30 hydrogen; then if h is absent and i is present, P_6 must be present and cannot be hydrogen;

- 111 -

vi) when P_1 and P_3 are oxo and b is present;
h, e, a, c and d are absent;
 P_2 = methyl; P_4 , P_5 , P_7 , P_8 , P_9 , P_{10} , P_{11} and
 P_{13} are hydrogen; then

- (a) if P_6 is absent, j cannot be a
double bond;
(b) if P_6 is hydrogen, i or j must be
present;

vii) when P_1 = OMethyl, a and c are present;
and P_2 = methyl; e, b, d and h are
absent and

P_3 , P_4 , P_7 , P_8 , P_9 , P_{10} , P_{11} and P_{13} are
hydrogen; then

- (a) if P_6 is hydrogen, i or j must be
present;
(b) if P_6 is absent, i and j cannot
both be double bonds;

viii) when P_1 = oxo and a, b, c, d, e are
absent;

P_2 = methyl; and

P_3 , P_4 , P_7 , P_9 , P_{10} , P_{11} , P_{12} and P_{13} are
hydrogen; then

- (a) if h is absent and j is present as
a double bond; P_6 cannot be
hydrogen;
(b) if h, i and j are absent, P_6 cannot
be hydrogen; and
(c) if h is present, j cannot be a
triple bond;

ix) when P_1 = OH, a, b, c, d, e and h are
absent;

P_2 = methyl;

- 112 -

$P_3, P_4, P_7, P_8, P_9, P_{10}, P_{11}, P_{12}$ and P_{13}
are hydrogen; then

- 5 (a) if j is a double bond and i is
absent; then P_6 is not hydrogen;
(b) if i is a double bond and j is
absent, then P_6 is not hydrogen;
(c) i and j cannot both be double
bonds;
10 (d) if i and j are absent, P_6 cannot be
hydrogen;

x) when $P_1 = \text{oxo}$, e and b are present; and
 a, c, d are absent;

$P_2 = \text{methyl}$; and

15 $P_3, P_4, P_7, P_9, P_{10}, P_{11}$ and P_{13} are
hydrogen; then

- (a) if h is present, j cannot be a
triple bond;
(b) if h is absent and i is a double
bond, then P_6 is not hydrogen;
20 (c) if h is absent and i is a double
bond, then P_6 is not hydrogen;
(d) if h is absent and neither i nor j
is a double bond, then P_6 is not
hydrogen;

25 xi) when $P_1 = \text{oxo}$, $P_3 = \text{OH}$, b is present, a ,
 e, c, d, h, i, j are absent, $P_2 = \text{methyl}$
and $P_4, P_7, P_9, P_{10}, P_{11}$ and P_{13} are
hydrogen; then P_6 cannot be hydrogen;

- 113 -

- xii) when $P_1 = \text{oxo}$, b is present, a, e, c, d, h, i are absent, P_6 and P_2 are methyl and

$P_3, P_4, P_5, P_7, P_8, P_9, P_{10}$ and P_{11} are hydrogen; then

- (a) if j is a double bond, P_{13} cannot be ethylenyl;
(b) if neither j nor k is a double or triple bond, then P_{13} cannot be acetylenyl;

- xiii) when $P_1 = \text{OH}$, c is present and a, b, e, d, h and i are absent;

P_2 and P_6 are methyl; and

$P_3, P_4, P_5, P_7, P_8, P_9, P_{10}, P_{11}$ and P_{12} are hydrogen; then

- (a) if j is a double bond, P_{13} is not ethyl;
(b) if j is absent and k is a double bond, P_{13} is not methyl-methylenyl;
(c) if j and k are absent; P_{13} is not ethyl; and
(d) if k is a triple bond, P_{13} cannot be methyl-methinyl.

1/96




FIG. 1A.




FIG. 1B.






FIG. 1C.


2/96

**FIG. 2.**


3/96

**FIG. 3.**

4/96

**FIG. 4.**

5/96

**FIG. 5.**

6/96

PREGNA-4, 16-DIEN-3-ONE





FIG. 6.


7/96

PREGNA-4, 16-DIEN-3-ONE

**FIG. 7.**


8/96

PREGNA-4, 16-DIEN-3-ONE

**FIG. 8.**

9/96

PREGNA-4, 16-DIEN - 3-ONE

**FIG. 9.**

10/96

PREGNA-4, 16-DIEN - 3 - ONE





FIG. 10.

11/96

PREGNA-5, 16-DIEN-3 β -OL**FIG. II.**

10/96

PREGNA-4, 16-DIEN - 3 - ONE






FIG. 10.

11/96

PREGNA-5, 16-DIEN-3 β -OL**FIG. II.**

12/96

PREGNA-5, 16-DIEN-3 β -OL*FIG. 12.*

13/96






FIG. 13.


14/93

**FIG. 14.**


15/96

PREGNA-5, 16-DIEN-3 β -OL**FIG. 15.**


16/96

PREGNA-5, 16-DIEN-3 β -OL**FIG. 16.**


17/96

**FIG. 17.**

18/96

*FIG. 18.*

19/96

**FIG. 19.**

20/96






FIG. 20.


21/96

*FIG. 21.*


22/96

*FIG. 22.*


23/96

*FIG 23.*


24/96

**FIG. 24.**


25/96

**FIG. 25.**


26/96

**FIG. 26.**


27/96

*FIG. 27.*


28/96

**FIG. 28.**

29/96

*FIG. 29.*

30/96

**FIG. 30.**

31/96

20, 21-DIMETHYL-PREGNA-5, 20-DIEN-3 β -OL**FIG. 31.**

32/96






FIG. 32.

33/96

**FIG. 33.**

34/96

**FIG 34.**

35/96




FIG. 35.

36/96

20, 21-DIMETHYL-PREGNA-5, 20-DIEN-3-ONE




FIG. 36.

37/96

20, 21-DIMETHYL-PREGNA-5, 20-DIEN-3-ONE




FIG. 37.

38/96





FIG. 38.

39/96

20, 21-DIMETHYL-PREGNA-5, 20-DIEN-3-ONE

**FIG. 39.**

40/96

20, 21-DIMETHYL-PREGNA-5, 20-DIEN-3-ONE




FIG. 40.

41/96


 $6\beta, 19$ -EPOXY-PREGNA-4, 17-DIEN-3-ONE

FIG. 41.

42/96


 $6\beta,19$ -EPOXY-PREGNA-4,17-DIEN-3-ONE

FIG. 42.

43/96




FIG. 43.

44/96


 $6\beta,19$ -EPOXY-PREGNA-4,17-DIEN-3-ONE

FIG. 44.

45/96


 $6\beta, 19$ -EPOXY-PREGNA-4, 17-DIEN-3-ONE

FIG. 45.

46/96




FIG. 46.

47/96





FIG. 47.

48/96

*FIG. 48.*

49/96




FIG. 49.

50/96





FIG. 50.

51/96

*FIG. 51.*

52/96





FIG 52.

53/96

*FIG. 53.*

54/96






FIG. 54.


55/96

*FIG. 55.*

56/96

**FIG 56.**

57/96

*FIG. 57.*

58/96






FIG. 58.


59/96

**FIG. 59.**


60/96

**FIG. 60.**


61/96

**FIG. 61.**


62/96

**FIG. 62.**


63/96

**FIG. 63.**


64/96

**FIG 64.**


65/96

*FIG. 65.*


66/96

**FIG. 66.**


67/96

**FIG. 67.**


68/96

**FIG. 68.**


69/96

*FIG. 69.*


70/96

**FIG. 70.**


71/96

**FIG 71.**


72/96

**FIG. 72.**


73/96

**FIG. 73.**


74/96

**FIG. 74.**

75/96

**FIG. 75.**

76/96

**FIG. 76.**

77/96




FIG. 77.

78/96




FIG. 78.

79/96






FIG. 79.


80/96

*FIG. 80.*


81/96

**FIG. 81.**


82/96

*FIG. 82.*


83/96

*FIG. 83.*


84/96

**FIG. 84.**


85/96

*FIG. 85.*


86/96

*FIG. 86.*


87/96

*FIG. 87.*


88/96

*FIG. 88.*


89/96

*FIG. 89.*


90/96

*FIG. 90.*

91/96

**FIG. 91.**

92/96

*FIG. 92.*

93/96






FIG. 93.


94/96

*FIG. 94.*

95/96

*FIG. 95.*

96/96

**FIG. 96.**

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/09634

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : CO7I 7/00, 9/00, 13/00 53/00 71/00; A61K 31/56, 31/57, 31/575, 31/58 US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC																				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/172, 177, 178, 181, 182; 540/48, 76, 91; 552/530, 532, 533, 551, 552, 553, 554, 555, 582, 603, 604, 605, 606, 607, 608, 609 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)																				
C. DOCUMENTS CONSIDERED TO BE RELEVANT																				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																		
A,P	WO, A, 94/28904 (PHERIN CORPORATION) 22 DECEMBER 1994. SEE ENTIRE DOCUMENT.	1-32																		
A	TETRAHEDRON, VOL 42, NO 2, ISSUED 1986, TERASAWA, ET AL., "CONVENIENT PREPARATIVE ROUTES TO 19-HYDROXY, 19-OXO-, 19-OIC-, AND 19-NOR-DEOXYCORTICOSTERONE," pages 537-545.	33																		
A	STEROIDS, VOL 59, ISSUED AUGUST 1994, LI, ET AL., "TWO POLYHYDOXYLATED STEROIDS FROM THE CHINESE SOFT CORAL LITOPHYTON ARBOREUM," pages 503-505.	33																		
A,P	WO, A, 94/28903 (PHERIN CORPORATION) 22 DECEMBER 1994, SEE ENTIRE DOCUMENT.	1-32																		
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.																				
<table border="0"><tr><td>* Special categories of cited documents:</td><td>"T"</td><td>later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principles or theory underlying the invention</td></tr><tr><td>"A" document defining the general state of the art which is not considered to be of particular relevance</td><td>"X"</td><td>document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td></tr><tr><td>"E" earlier document published on or after the international filing date</td><td>"Y"</td><td>document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td></tr><tr><td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td><td>"A"</td><td>document member of the same patent family</td></tr><tr><td>"O" document referring to an oral disclosure, use, exhibition or other means</td><td></td><td></td></tr><tr><td>"P" document published prior to the international filing date but later than the priority date claimed</td><td></td><td></td></tr></table>			* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principles or theory underlying the invention	"A" document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"E" earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A"	document member of the same patent family	"O" document referring to an oral disclosure, use, exhibition or other means			"P" document published prior to the international filing date but later than the priority date claimed		
* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principles or theory underlying the invention																		
"A" document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																		
"E" earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																		
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A"	document member of the same patent family																		
"O" document referring to an oral disclosure, use, exhibition or other means																				
"P" document published prior to the international filing date but later than the priority date claimed																				
Date of the actual completion of the international search 18 OCTOBER 1995		Date of mailing of the international search report 03 NOV 1995																		
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer KIMBERLY J. KESTLER Telephone No. (703) 308-1235																		

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/09634

A. CLASSIFICATION OF SUBJECT MATTER: US CL :

514/172, 177, 178, 181, 182; 540/48, 76, 91; 552/530, 532, 533, 551, 552, 553, 554, 555, 582, 603, 604, 605, 606, 607, 608, 609

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.